



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 418 071 A2**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90310009.7

(22) Date of filing: 13.09.90

(51) Int. Cl.⁵: **C07C 323/60**, C07D 239/58,
C07D 213/75, C07D 213/70,
C07C 323/52, A61K 31/16,
A61K 31/33, C07D 307/64,
C07D 277/74, C07D 277/36,
C07D 215/36

(30) Priority: 15.09.89 PCT/US89/04033

(43) Date of publication of application:
20.03.91 Bulletin 91/12

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: **PFIZER INC.**
235 East 42nd Street
New York, N.Y. 10017(US)


(72) Inventor: **McCarthy, Peter A.**
235 North Anguilla
RR5, Pawcatuck, Connecticut(US)
Inventor: **Walker, Frederick J.**

5, Maynard Hill Road
Preston, Connecticut(US)
Inventor: **Truong, Thien**
14, Cinnamon Ridge
Old Saybrook, Connecticut 06475(US)
Inventor: **Hamanaka, Ernest S.**
40 Eagle Ridge Drive
Gales Ferry, Connecticut(US)
Inventor: **Chang, George**
One Winthrop Hill Road
Ivoryton, Connecticut 06442(US)

(74) Representative: **Moore, James William, Dr.**
Pfizer Limited Ramsgate Road
Sandwich Kent CT13 9NJ(GB)

(54) New N-aryl and N-heteroarylamide and urea derivatives as inhibitors of acyl coenzyme A: cholesterol acyl transferase.

(57) Compounds of the formula



the pharmaceutically acceptable salts thereof, wherein Q and R¹ are as defined below, and novel carboxylic acid and acid halide intermediates used in the synthesis of such compounds. The compounds of formula I are inhibitors of acyl coenzyme A: cholesterol acyltransferase (ACAT) and are useful as hypolipidemic and antiatherosclerosis agents.

EP 0 418 071 A2

NEW N-ARYL AND N-HETEROARYLAMIDE AND UREA DERIVATIVES AS INHIBITORS OF ACYL COENZYME A: CHOLESTEROL ACYL TRANSFERASE (ACAT)


The present invention relates to new N-aryl and N-heteroarylamide and urea derivatives, pharmaceutical compositions comprising such compounds, novel carboxylic acid and acid halide intermediates used in the synthesis of such compounds and the use of such compounds to inhibit intestinal absorption of cholesterol, lower serum cholesterol and reverse the development of atherosclerosis. The compounds are inhibitors of acyl coenzyme A: cholesterol acyltransferase (ACAT).

Cholesterol that is consumed in the diet (dietary cholesterol) is absorbed as free cholesterol by the mucosal cells of the small intestine. It is then esterified by the enzyme ACAT, packaged into particles known as chylomicrons, and released into the bloodstream. Chylomicrons are particles into which dietary cholesterol is packaged and transported in the bloodstream. By inhibiting the action of ACAT, the compounds of this invention prevent intestinal absorption of dietary cholesterol and thus lower serum cholesterol levels. They are therefore useful in preventing atherosclerosis, heart attacks and strokes.

By inhibiting the action of ACAT, the compounds of the present invention also enable cholesterol to be removed from the walls of blood vessels. This activity renders such compounds useful in slowing or reversing the development of atherosclerosis as well as in preventing heart attacks and strokes.





Other inhibitors of ACAT are referred to in United States Patents 4,716,175 and 4,743,605 (a divisional of the '175 patent) and in the European Patent Applications having publication numbers 0 242 610, 0 245 687 and 0 252 524. Certain ureas and thioureas as antiatherosclerosis agents are referred to in United States Patent 4,623,662.

The present invention relates to compounds of the formula



wherein Q is -CR²R³R⁴ or -NR¹⁷R¹⁸:

R¹ is



R², R³ and R⁴ may be the same or different, and

(a) are selected from the group consisting of hydrogen, (C₁-C₄) alkyl, A, XR¹⁰, phenyl-(C₁-C₇) alkyl, and (C₅-C₆) cycloalkyl-(C₁-C₆) alkyl, with the proviso that at least one of R², R³ and R⁴ must be A, and with the proviso that when R¹ is a group of the formula XXVII or a group of the formula XXVI wherein G is nitrogen and wherein neither R⁵, R⁶ nor R¹⁵ is NR¹⁹R²⁰, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄) alkylthio, phenylthio or heteroalkylthio, either at least one of R², R³ and R⁴ must be XR¹⁰, or two of R², R³ and R⁴ must be A; or

(b) R² and R³ together with the carbon to which they are attached form a cyclic or bicyclic system selected from the group consisting of (C₃-C₇) cycloalkyl, (C₃-C₇) cycloalkenyl, (C₆-C₁₄) bicycloalkyl, (C₆-C₁₄) bicycloalkenyl, and aryl-fused and heteroaryl-fused systems containing 8 to 15 carbon atoms, one ring of any of said aryl-fused and heteroaryl-fused systems being aromatic and the ring containing the carbon to which R² and R³ are attached being non-aromatic, one of the carbons of said aromatic ring being optionally replaced by sulfur or oxygen, one or more carbons of said non-aromatic ring being optionally replaced by sulfur or oxygen, and one or more carbons of said aromatic ring being optionally replaced by nitrogen; one or two carbons of said cycloalkyl or bicycloalkyl groups being optionally replaced by sulfur or oxygen, and said cyclic or bicyclic system being optionally substituted with one to five substituents independently selected from the group consisting of phenyl, substituted phenyl, (C₁-C₆) alkyl and A, with the proviso that one and only one of said substituents is A, and one and only one of said substituents is phenyl or substituted phenyl, said substituted phenyl being substituted with one or more substituents independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) alkylthio, halogen and trifluoromethyl; and R⁴ is hydrogen, XR¹⁰ or A;

with the proviso that when R¹ is a group of the formula XXVII or a group of the formula XXVI wherein G is nitrogen and wherein neither R⁵, R⁶ nor R¹⁵ is NR¹⁹R²⁰, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄) alkylthio, phenylthio or heteroalkylthio, R² and R³, together with the carbon to which they are attached, do not form a (C₃-C₇) cycloalkyl ring containing only carbon atoms:

A is a hydrocarbon containing 4 to 16 carbons and 0, 1 or 2 double bonds:

X is O, S, SO, SO₂, NH, NR²³CO or NSO₂R²⁴, wherein R²³ is hydrogen or (C₁-C₆) alkyl and R²⁴ is (C₁-C₆) alkyl, phenyl or (C₁-C₃) alkyl-phenyl;

R⁵, R⁶, R¹⁵ and R¹⁶ are each independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, iodo, (C₁-C₄) alkyl, (C₁-C₄) haloalkyl, (C₁-C₄) alkoxy, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄) alkylthio, substituted phenylthio, heteroarylthio, heteroaryloxy, and NR¹⁹R²⁰, wherein R¹⁹ and R²⁰ are the same or different and are selected from the group consisting of hydrogen, (C₁-C₄) alkyl, phenyl, substituted phenyl, (C₁-C₄) acyl, aroyl, and substituted aroyl, wherein said substituted phenyl and substituted aroyl groups are substituted with one or more substituents independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) alkylthio, halogen and trifluoromethyl, or R¹⁹ and R²⁰, together with the nitrogen to which they are attached, form a piperidine or morpholine ring; and wherein R⁵, R⁶, R¹⁵ and R¹⁶, when attached to a bicyclic system, may be attached to either ring of such system, with the proviso that no more than 3 non-hydrogen substituents may be attached to any one ring of such system:

R⁷, R⁸ and R⁹ are the same or different; R⁷ is selected from the group consisting of hydrogen, (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, methyl and fluoro; and R⁸ and R⁹ are each independently selected from the group consisting of (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, methyl, and fluoro:

R¹⁰ is selected from the group consisting of (C₄-C₁₂) cycloalkyl, (C₄-C₁₂) straight or branched alkyl, (C₄-C₁₂) cycloalkyl-(C₁-C₆) alkyl, phenyl-(C₁-C₆) alkyl, (substituted phenyl)-(C₁-C₆) alkyl, (C₁-C₆) alkyl-phenyl, (C₁-C₆) alkyl-(substituted phenyl), optionally substituted thiazoles, optionally substituted benzothiazoles, and optionally substituted pyridines; wherein the substituents on the substituted phenyl, substituted thiazoles, substituted benzothiazoles and substituted pyridines are selected from the group consisting of (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, (C₁-C₆) alkyl, halo and trifluoromethyl:

B, D, E and G are selected from the group consisting of nitrogen and carbon, with the proviso that one or more of B, D and E is nitrogen, and with the proviso that when G is nitrogen, the group XXVI is attached to the nitrogen of formula I at the 4 or 5 position of the pyrimidine ring (designated by a and b):

and R¹⁷ and R¹⁸ are each independently selected from the group consisting of (C₄-C₁₂) straight or branched alkyl, phenyl-(C₁-C₄) alkyl, and (C₁-C₆) alkylphenyl-(C₁-C₆) alkyl:

with the proviso that when Q is NR¹⁷R¹⁸, R¹ is a group of the formula XXVI or XXIV, or a group of the formula XXVII wherein R⁷, R⁸ and R⁹ are each methoxy.

Examples of said aryl-fused and heteroaryl-fused systems are:

1,2,3,4-tetrahydronaphthalene,

5,6,7,8,9-pentahydrobenzocycloheptene,

- 5,6,7,8,9,10-hexahydrobenzocyclooctene,
 4,5,6-trihydro-1-thiapentalene,
 4,5,6-trihydro-2-thiapentalene,
 4,5,6,7-tetrahydrobenzo[b]thiophene,
 5 4,5,6,7-tetrahydrobenzo[c]thiophene,
 4,5,6-trihydro-1-oxapentalene,
 4,5,6,7-tetrahydrobenzo[b]furan,
 4,5,6-trihydro-1-azapentalene,
 4,5,6,7-tetrahydrobenzo[b]pyrrole,
 10 4,5,6-trihydro-1-oxa-3-azapentalene,
 4,5,6,7-tetrahydrobenzo[d]oxazole,
 4,5,6-trihydro-1-thia-3-azapentalene,
 4,5,6,7-tetrahydrobenzo[d]thiazole,
 4,5,6-trihydro-1-oxa-2-azapentalene,
 15 4,5,6,7-tetrahydrobenzo[d]oxazole,
 4,5,6-trihydro-1-thia-2-azapentalene,
 4,5,6,7-tetrahydrobenzo[d]thiazole,
 4,5,6-trihydro-1,2-diazapentalene,
 4,5,6,7-tetrahydrobenzo[d]pyrazole,
 20 4,6-diazaindane and
 5,6,7,8-tetrahydroquinazoline.

Unless otherwise indicated, the term "halo", as used herein, includes fluoro, chloro, bromo and iodo.

Unless otherwise indicated, the term "alkyl", as used herein, may be straight, branched or cyclic, and may include straight and cyclic moieties as well as branched and cyclic moieties.


- 25 Unless otherwise indicated, the term "one or more substituents", as used herein, refers to from one to the maximum number of substituents possible based on the number of available bonding sites.

The term "one or more carbons of said non-aromatic ring", as used herein, refers to from one to all of the carbon atoms that are part of the non-aromatic ring of any of the aryl-fused or heteroaryl-fused systems described above, and not part of the aromatic ring of said aryl-fused system.

- 30 The term "one or more carbons of said aromatic ring", as used herein, refers to from one to all of the carbon atoms that are part of the aromatic ring of any of the aryl-fused and heteroaryl-fused systems described above, or are part of both said aromatic and non-aromatic rings of said aryl-fused and heteroaryl-fused system.

- The compounds of formula I may have optical centers and therefore may occur in different stereoisomeric configurations. The invention includes all stereoisomers of such compounds of formula I, including mixtures thereof.


The present invention also relates to compounds of the formula



- 45 wherein Z is hydroxy, chloro or bromo; R², R³ and R⁴ are each independently selected from the group consisting of (C₆-C₁₆) alkyl, (C₆-C₁₆) alkenyl containing 1 or 2 double bonds, phenyl-(C₁-C₆) alkyl, (C₅-C₆) cycloalkyl-(C₁-C₆) alkyl, XR¹⁰, and hydrogen, with the proviso that at least one of R², R³ and R⁴ is (C₆-C₁₆) alkyl or (C₆-C₁₆) alkenyl containing 1 or 2 double bonds; or R² and R³, together with the carbon to which they are attached, form a cyclic or bicyclic ring system selected from the group consisting of

50


55



and aryl-fused and heteroaryl-fused systems containing 8 to 15 carbon atoms, one ring of any of said aryl-fused and heteroaryl-fused systems being aromatic and the other ring containing the carbon to which R^2 and R^3 are attached being non-aromatic, one of the carbons of said aromatic ring being optionally replaced by sulfur or oxygen, one or more carbons of said non-aromatic ring being optionally replaced by sulfur or oxygen, and one or more carbons of said aromatic ring being optionally replaced by nitrogen; and wherein R^4 is hydrogen, A or XR^{10} ; A is a hydrocarbon containing 4 to 16 carbons and 0, 1 or 2 double bonds; X is oxygen or sulfur; R^{10} is selected from the group consisting of (C_4-C_{12}) cycloalkyl, (C_4-C_{12}) straight or branched alkyl, (C_4-C_{12}) cycloalkyl- (C_1-C_6) alkyl, phenyl- (C_1-C_6) alkyl, (substituted phenyl)- (C_1-C_6) alkyl, (C_1-C_6) alkyl-phenyl, (C_1-C_6) alkyl-(substituted phenyl), substituted thiazoles, substituted benzothiazoles, and substituted pyridines, wherein the substituents on the substituted phenyl, substituted thiazoles, substituted benzothiazoles and substituted pyridines are selected from the group consisting of (C_1-C_4) alkoxy, (C_1-C_4) alkylthio, (C_1-C_6) alkyl, halo and trifluoromethyl; R^{11} is (C_6-C_{12}) alkyl or (C_6-C_{12}) alkenyl containing 1 or 2 double bonds; n and m are each independently 1 or 2; o and p are each independently 0, 1 or 2; each broken line represents an optional double bond; and each asterisk represents the carbon to which R^2 and R^3 are attached; with the proviso that when R^2 and R^3 form any of ring systems XVIII, XIX and XX, R^4 is A or XR^{10} , and when R^2 and R^3 form any of ring systems XXI, XXII and XXIII, R^4 is hydrogen; and with the proviso that when R^2 and R^3 do not form a ring system, no more than one of R^2 , R^3 and R^4 is hydrogen. The compounds of formula II are intermediates used in the synthesis of compounds of the formula I.

Specific compounds of the formula II are: 2-(hexylthio)octanoic acid, 2-(hexylthio)nonanoic acid, 2-(hexylthio)decanoic acid, 2-(heptylthio)octanoic acid, 2-(heptylthio)nonanoic acid, 2-(heptylthio)decanoic acid, 2-nonylindan-2-yl carboxylic acid, 2-decylindan-2-yl carboxylic acid, 2-octyl-1,2,3,4-tetrahydronaphth-2-yl carboxylic acid and 2-nonyl-1,2,3,4-tetrahydronaphth-2-yl carboxylic acid.

The present invention also relates to compounds of the formula



XXVIII

5

wherein R²¹ is (C₁-C₃) alkyl and R²² is hydrogen or (C₁-C₃) alkyl.

- 10 Preferred compounds of formula I are those wherein R¹ is 2,4,6-trifluorophenyl, 2,4,6-trimethoxyphenyl, 6-methoxy quinolin-5-yl, 6-methylthioquinolin-5-yl, 6-methoxy-isoquinolin-5-yl, 6-methylthioisoquinolin-5-yl, 6-methylthio-8-acetaminoquinolin-5-yl, 2-methyl-4,6-di(methylthio)pyrimidin-5-yl, and 6-methyl-2,4-di(methylthio)pyridin-3-yl.

Other preferred compounds of formula I are those wherein:

- 15 R² is hexylthio, R³ is octyl and R⁴ is hydrogen; or
R² and R³ together with the carbon to which they are attached form an Indan-2-yl ring, and R⁴ is 2-decyl; or
R² and R³ together with the carbon to which they are attached form a 1,2,3,4-tetrahydronaphth-2-yl ring and R⁴ is nonyl.

Specific preferred compounds of formula I are:

- 20 N-(2,4,6-trifluorophenyl)-2-(hexylthio) octanoic amide;
N-(2,4,6-trimethoxyphenyl)-2-(hexylthio) octanoic amide;
N-(2,4,6-trimethoxyphenyl)-2-(6-ethoxybenzothiazol-2-yl) decanoic amide;
N-(6-methoxyquinolin-5-yl)-2-(hexylthio) decanoic amide;
N-(6-methylthioquinolin-5-yl)-2-(hexylthio) decanoic amide;
25 N-(6-methoxyisoquinolin-5-yl)-2-(hexylthio) decanoic amide;
N-(6-methylthio-8-acetaminoquinolin-5-yl)-2-(hexylthio)decanoic amide;
N-(6-methylthioisoquinolin-5-yl)-2-(hexylthio) decanoic amide;
N-(6-methylisoquinolin-5-yl)-2-(hexylthio) decanoic amide;
N-(6-methylthioquinolin-5-yl)-2-(4-(3-methylpropyl)phenoxy) nonanoic amide;
30 (2S)-N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-hexylthiodecanoic amide;
(2S)-N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide;
(2S)-N-[6-(methylthio)quinolin-5-yl]-2-hexylthiodecanoic amide;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-2-hexylthiodecanoic amide;
N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-hexylthiodecanoic amide;
35 N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-2,2-dimethyldodecanoic amide;
N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2,2-dimethyldodecanoic amide;
N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-N-[4-(3-methylbutyl)benzyl]-N-cycloheptylurea;
N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-N-[4-(3-methylbutyl)benzyl]-N-heptylurea;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-N-[4-(3-methylbutyl)benzyl]-N-cycloheptylurea;
40 N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-N-[4-(3-methylbutyl)benzyl]-N-heptylurea;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-4,5-dimethyl-trans-2-heptylcyclohex-4-ene-carboxamide;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-2-heptylnonanoic amide;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]pentadecanoic amide;
N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]pentadecanoic amide;
45 N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-(Z)-9-octadecenoic amide;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-(Z)-9-octadecenoic amide;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-trans-3-nonyl-1,2,3,4-tetrahydro-2-naphthoic amide;
N-[4,6-bis(methylthio)pyrimidin-5-yl]-trans-3-nonyl-1,2,3,4-tetrahydro-2-naphthoic amide;
N-(2,4,6-trimethoxyphenyl)-2-nonyl-1,2,3,4-tetrahydronaph-2-yl carboxamide;
50 N-(2,4,6-trifluorophenyl)-2-nonyl-1,2,3,4-tetrahydronaph-2-yl carboxamide;
N-(2,4,6-trifluorophenyl)-2-nonylindan-2-yl carboxamide;
N-(2,4,6-trimethoxyphenyl)-6-trans-heptyl-3,4-dimethylcyclohex-2-enyl carboxamide;
N-(2,4,6-trimethoxyphenyl)-2-nonylbicyclo[2.2.1]hept-5-en-2-yl carboxamide;
N-(6-methylthioquinolin-5-yl)-2-octyl-1,3-dithian-2-yl carboxamide;
55 N-(2-methyl-4,6-dimethylthiopyrimidin-5-yl)-2-hexylthiodecanoic amide;
N-(2,4,6-trimethoxyphenyl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea;
N-(2,4,6-trimethoxyphenyl)-N'-(4-(2,2-dimethylpropyl)phenylmethyl)-N'-heptylurea;
N-(isoquinolin-5-yl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea;

N-(quinolin-5-yl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea; and
 N-(6-methoxyquinolin-5-yl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea.

Other compounds of formula I are:

- N-(2-methoxy-6-methylphenyl)-2-(4-propylphenyl)sulfonylundecanoic amide;
- 5 N-(3-methyl-6-trifluoromethylquinolin-5-yl)-2-(2,5-dimethylphenoxy)decanoic amide;
- N-(6,7-(methylenedioxy)isoquinolin-5-yl)-2-(thiazol-2-yl)methoxynonanoic amide;
- N-(2,6-dimethoxyphenyl)-2,2-di(isopropylthio)octanoic amide;
- N-(3-chloro-8-isobutylquinolin-5-yl)-2-(3-ethoxyphenyl)thio-2-propylheptanoic amide;
- N-(4-methyl-6,7-difluoroisoquinolin-5-yl)-1-methyl-5-octyl-2,7-dioxabicyclo[2.2.1]heptan-5-yl carboxamide;
- 10 N-(2,4,6-trimethoxyphenyl)-2-propylthiohexadec-9-enoic amide;
- N-(6-isopropylthioquinolin-5-yl)-2-(4-ethylthiophenoxy)nonanoic amide;
- N-(6-methoxyisoquinolin-5-yl)-2-[5-chlorobenzthiazol-2-yl]thio]octanoic amide;
- N-(2,4,6-trifluorophenyl)-2-[(3,5-dimethylbenzoyl)amino]octanoic amide;
- N-(8-acetamino-6-methoxyquinolin-5-yl)-2-hexylthiodecanoic amide;
- 15 N-(8-amino-6-methoxyquinolin-5-yl)-2-hexylthiodecanoic amide;
- N-(8-amino-6-methylthioquinolin-5-yl)-2-hexylthiodecanoic amide;
- N-(8-(2,2-dimethylpropionyl)amino-6-methoxyquinolin-5-yl)-2-hexylthiodecanoic amide;
- N-(8-(2,2-dimethylpropionyl)amino-6-methoxyquinolin-5-yl)-2-hexylthiodecanoic amide;
- N-(6-methylthioquinolin-5-yl)-2-hexyloxydecanoic amide;
- 20 N-(8-acetamino-6-methylthioquinolin-5-yl)-2-hexyloxydecanoic amide;
- N-(6-methylthioquinolin-5-yl)-2-heptylnonanoic amide;
- N-(8-acetamino-6-methylthioquinolin-5-yl)-2-heptylnonanoic amide;
- N-(8-acetaminoquinolin-5-yl)-2-hexylthiodecanoic amide;
- N-(8-aminoquinolin-5-yl)-2-hexylthiodecanoic amide;
- 25 N-(8-acetaminoquinolin-5-yl)-2-hexylthiodecanoic amide;
- N-(6-aminoquinolin-5-yl)-2-hexylthiodecanoic amide;
- N-(4,6-dimethylthiopyrimidin-5-yl)-2-hexylthiodecanoic amide;
- N-(4,6-diethylthiopyrimidin-5-yl)-2-hexylthiodecanoic amide;
- N-(4-methoxy-6-ethylthiopyrimidin-5-yl)-2-hexylthiodecanoic amide;
- 30 N-(4-ethoxy-6-methylthiopyrimidin-5-yl)-2-hexylthiodecanoic amide;
- N-(4-ethoxy-6-ethylthiopyrimidin-5-yl)-2-hexylthiodecanoic amide;
- N-(4-methoxy-6-ethoxyethylthiopyrimidin-5-yl)-2-hexylthiodecanoic amide;
- N-(4-methoxy-6-butylthiopyrimidin-5-yl)-2-hexylthiodecanoic amide;
- N-(2,4-dimethylthio-6-methylpyrimidin-5-yl)-2-heptylnonanoic amide;
- 35 N-(2-amino-4-methoxy-6-methylthiopyrimidin-5-yl)-2-hexylthiodecanoic amide;
- N-(2-acetamino-4-methoxy-6-methylthiopyrimidin-5-yl)-2-hexylthiodecanoic amide;
- N-[4-methoxy-6-(2-furylmethylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide;
- N-[4-methoxy-6-(2-propylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide;
- N-(2-butylthio-4-methylpyridin-3-yl)-2-hexylthiodecanoic amide;
- 40 N-[2-(4-methoxyphenylthio)-4-methylpyridin-3-yl]-2-hexylthiodecanoic amide;
- N-[2-(2-furylmethylthio)-4-methylpyridin-3-yl]-2-hexylthiodecanoic amide;
- N-(2-ethylthio-4-methylpyridin-3-yl)-2-hexylthiodecanoic amide;
- N-(6-methoxyquinolin-5-yl)-9-octadecenoic amide;
- N-(6-fluoroquinolin-5-yl)-9-octadecenoic amide;
- 45 N-(6-methylthioquinolin-5-yl)-9-octadecenoic amide;
- N-(8-acetamino-6-methylthio-quinolin-5-yl)-9-octadecanoic amide;
- N-(2-ethylthio-4-methylpyridin-3-yl)-4,5-dimethyl-trans-2-nonylcyclohex-4-enecarboxamide; and
- N-(4,6-dimethylthiopyrimidin-5-yl)-2-decylindane-2-carboxamide.

The present invention also relates to all radiolabelled forms of the compounds of the formulae I, II and
 50 XXVIII. Such radiolabelled compounds are useful as research and diagnostic tools in metabolism pharmacokinetic studies and in binding assays in both animals and man.

The present invention also relates to a pharmaceutical composition for inhibiting ACAT, inhibiting intestinal absorption of cholesterol, reversing or slowing the development of atherosclerosis, or lowering the concentration of serum cholesterol in a mammal, including a human, comprising an amount of a compound
 55 of the formula I, or a pharmaceutically acceptable salt thereof, effective in inhibiting ACAT, inhibiting intestinal absorption of cholesterol, reversing or slowing the development of atherosclerosis, or lowering the concentration of serum cholesterol, and a pharmaceutically acceptable carrier.

The present invention also relates to a method for inhibiting ACAT, inhibiting intestinal absorption of


cholesterol, reversing or slowing the development of atherosclerosis, or lowering the concentration of serum cholesterol in a mammal, including a human, comprising administering to a mammal an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in inhibiting ACAT, inhibiting intestinal absorption of cholesterol, reversing or slowing the development of atherosclerosis, or lowering the concentration of serum cholesterol.

Examples of pharmaceutically acceptable acid addition salts of the compounds of formula I salts are the salts of hydrochloric acid, p-toluenesulfonic acid, fumaric acid, citric acid, succinic acid, salicylic acid, oxalic acid, hydrobromic acid, phosphoric acid, methanesulfonic acid, tartaric acid, di-p-toluoyl tartaric acid, and mandelic acid.


Reaction schemes 1-4 below illustrate the synthesis of the compounds of this invention. The compounds of formula I designated in the reaction schemes by the formulae IA, IB, IC, and ID, depending on the method by which they prepared. Scheme 5 below illustrates the synthesis of certain 5-aminoquinolines and 5-aminoisoquinolines used as reactants in scheme 1-4.



Except where otherwise stated, Q, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹⁵, R¹⁶, R¹⁷, R¹⁸, n, m, o, p, X, A, B, D, and G in the reaction schemes and discussion that follows are defined as above.


SCHEME 1



SCHEME 2



SCHEME 3SCHEME 4

SCHEME 5

Scheme 1 represents the synthesis of amides of the present invention having the formula IA, i.e., compounds of the formula I wherein Q is $-\text{CR}^2\text{R}^3\text{R}^4$, from the corresponding carboxylic acid having the formula III. An acid of formula III is first converted to the corresponding acid halide of formula IV, wherein W is chloro or bromo, by reacting it with a chlorinating or brominating agent. Examples of suitable chlorinating and brominating agents are oxalyl chloride, oxalyl bromide, thionyl chloride, thionyl bromide, phosphorous trichloride, phosphorous tribromide, phosphorous pentachloride, phosphorous pentabromide, phosphorous oxychloride, and phosphorous oxybromide. This reaction is typically carried out in the absence of a solvent or, alternatively, in the presence of a halogenated hydrocarbon solvent such as methylene chloride, for from about .5 to 48 hours (preferably from about 2 to 18 hours) at a temperature from about 0-250°C (preferably at the reflux temperature of the reaction mixture). The acid halide so formed is then converted to the


corresponding amide of the formula IA by reacting it with an amine of the formula R^1NH_2 and an acid scavenger such as dimethylaminopyridine, pyridine or triethylamine. This reaction is typically carried out in the absence of a solvent or in the presence of an inert solvent such as tetrahydrofuran or methylene chloride for from about 0.25 to 144 hours (preferably from about 2 to 72 hours) at a temperature from about

5 -78 to 350 °C (preferably from about -20 to the reflux temperature of the reaction mixture).
 Compounds of the formula I, wherein Q is $-CR^2R^3R^4$, R^2 is XR^{10} , one of R^3 and R^4 is hydrogen and the other is selected from hydrogen, (C_1-C_4) alkyl or A, i.e., compounds of the formula IB, may be prepared as illustrated in scheme 2. Referring to scheme 2, a carboxylic acid of the formula V, wherein one of R^3 and R^4 is hydrogen and the other is selected from hydrogen, (C_1-C_4) alkyl or A, is reacted for about 3 hours
 10 with thionyl chloride using no solvent, at the reflux temperature of the reaction mixture. Bromine and a catalytic amount of iodine are then added to the reaction mixture, and the resulting mixture is brought to reflux. After refluxing for about 18 hours, ethanol is added and the mixture is refluxed for about 1 more hour to produce a bromoester of the formula VI, wherein R^3 and R^4 are defined as they are for formula V above. The bromoester of formula VI is then converted to an ester having the same formula as formula VI except
 15 that the substituent -Br is replaced by the substituent $-XR^{10}$, (hereinafter referred to as formula VI'), by reacting it with a compound of the formula HXR^{10} and a base such as potassium carbonate or sodium hydride in an aprotic, polar solvent such as dimethylformamide, acetone or tetrahydrofuran, for about .5 to 48 hours (preferably from about 4 to 18 hours) at a temperature from about -78 to 350 °C (preferably from about 0 °C to the reflux temperature of the reaction mixture). An acid of the formula VII, wherein R^3 and R^4
 20 are defined as they are for formulas V and VI above, is then prepared by reacting the ester having formula VI' with a hydroxide such as sodium hydroxide. This reaction is typically carried out overnight in an lower alcohol solvent such as methanol or ethanol, at a temperature from about -78 to 350 °C (preferably from about 20 °C to the reflux temperature of the reaction mixture).

The acid of formula VII so prepared is then converted to an amide of the formula IB, wherein R^3 and R^4
 25 are defined as they are for formulae V, VI and VII above, by the acid to amide synthesis illustrated in scheme 1 and described above.

Compounds of the formula IB, may be prepared, alternatively, by the following method. A compound of the formula V, as illustrated in scheme 2 and defined above, is reacted with thionyl chloride followed by bromine and a catalytic amount of iodine as described above, but quenching the reaction with water instead
 30 of ethanol, to form a compound of the formula $HOCCBrR^3R^4$, wherein R^3 and R^4 are defined as they are for formula V. This compound is then converted, sequentially, to the corresponding acid chloride of the formula $ClCOCBrR^3R^4$ and the corresponding amide of the formula $R^1NHCOCBrR^3R^4$, wherein R^3 and R^4 are defined as they are for formula V, by the acid to amide synthesis illustrated in scheme 1 and described above. The amide of the formula $R^1NHCOCBrR^3R^4$ so formed is then reacted with a compound of the
 35 formula HXR^{10} and a base such as potassium carbonate and sodium hydride to form a compound having the formula IB, wherein R^3 and R^4 are defined as they are for formula V. This reaction is typically carried out in an aprotic, polar solvent such as dimethylformamide, acetone or tetrahydrofuran, for from about 0.5 to 48 hours (preferably from about 4 to 18 hours). The reaction may be carried out at temperatures ranging from about -78 to 350 °C (preferably from about 0 °C to the reflux temperature of the reaction mixture).

40 Scheme 3 illustrates the preparation of compounds of formula I, wherein Q is $CR^2R^3R^4$, R^4 is hydrogen or A, and R^2 and R^3 , together with the carbon to which they are attached, form the bicyclic ring system



50 wherein the asterisk designates the carbon to which R^2 and R^3 are attached, and each of R^{12} and R^{13} are independently selected from the group consisting of hydrogen and (C_1-C_4) alkyl, or R^{12} and R^{13} , together with the carbons to which they are attached, form a benzene ring.

As illustrated in scheme 3, a Diels-Alder reaction is carried out between an acid of the formula XIII,
 55 wherein R^{11} is A, hydrogen, phenyl or substituted phenyl, and wherein R^{11} and the carboxyl group are trans to each other, and a diene of the formula IX, wherein R^{12} and R^{13} are as defined above. This reaction is typically carried out in a hydrocarbon solvent such as toluene, using a catalytic amount of an antioxidant such as hydroquinone. The reagents are generally reacted for about 1 to 10 days (preferably for about 3 to

5 days) in a sealed, high pressure apparatus at a temperature from about room temperature to 350° C (preferably from about 100 to 150° C). The reaction yields an acid of the formula X, which can be converted to the corresponding amide of the formula IC, wherein the carbons to which R¹² and R¹³ are attached are bonded by a carbon-carbon double bond, by the acid to amide synthesis illustrated in scheme 1 and described above. The amide of formula IC so formed can be converted to an amide of the formula IC, wherein the carbons to which R¹² and R¹³ are attached are bonded by a carbon-carbon single bond, by reacting it with a reducing agent such as hydrogen. Typically, the reduction is carried out using hydrogen gas in a high pressure apparatus, in an inert solvent such as acetic acid, and in the presence of a hydrogenation catalyst such as palladium on carbon.

10 The reduction may be carried out at temperatures ranging from about -20 to 250° C (preferably at room temperature). Fifty p.s.i. of hydrogen is the preferred pressure, through pressures greater than or equal to 1 atmosphere are suitable. The corresponding compound of formula IC, wherein the carboxyl group and R¹¹ are cis to each other, may be prepared in a similar manner, but using the corresponding cis isomer of the acid of formula XIII.

15 When the procedure of scheme 3 described above is used to prepare a compound of formula IC wherein R¹² and R¹³, together with the carbon to which they are attached, form a benzene ring, the diene of formula IX is generally generated in situ in the presence of the acid of formula XIII or its ester by heating a mixture of 1,3-dihydrobenzo[c]thiophene 2,2-dioxide and such acid or ester. This reaction is typically carried out at a temperature of from about 235 to 300° C (preferably from about 250 to 265° C) under nitrogen for approximately from 0.5 to 24 hours (preferably for about 2 hours).

20 Scheme 4 illustrates the preparation of compounds of the formula I, wherein Q is -CR²R³R⁴, R⁴ is (C₁-C₄) alkyl or A, and R² and R³ are each independently selected from the group consisting of hydrogen, (C₁-C₄) alkyl, A or XR¹⁰; or R² and R³, together with the carbon to which they are attached, form a cyclic or bicyclic system selected from the group consisting of (C₃-C₇) cycloalkyl, (C₆-C₁₄) bicycloalkyl, one or two carbons of said cycloalkyl and bicycloalkyl groups being optionally replaced with oxygen or sulfur; and aryl-fused and heteroaryl-fused systems containing 8 to 15 carbon atoms, one ring of any of said aryl-fused and heteroaryl-fused systems being aromatic and the ring containing the carbon to which R² and R³ are attached being non-aromatic, one of the carbons of said aromatic ring being optionally replaced by sulfur or oxygen, one or more carbons of said non-aromatic ring being optionally replaced by sulfur or oxygen, and one or more carbons of said aromatic ring being optionally replaced by nitrogen. It also illustrates the preparation of compounds of the formula I, wherein Q is -CR²R³R⁴, R⁴ is XR¹⁰, and R² and R³, together with the carbon to which they are attached, form a cyclic or bicyclic system as defined immediately above. All compounds of the invention illustrated in scheme 4 are designated by the formula ID and are prepared by the following procedure.

35 An acid of the formula XI, wherein R² and R³ are each independently selected from the group consisting of hydrogen, (C₁-C₄) alkyl, A or XR¹⁰, or R² and R³ together with the carbon to which they are attached, form a cyclic or bicyclic system as defined immediately above, is reacted with a base such as lithium diisopropylamide or hexamethyldisilazide, with or without an additive such as hexamethylphosphorous triamide, in a dry inert solvent such as tetrahydrofuran, and then reacted with a compound of the formula R⁴Hal, wherein Hal is halogen and R⁴ is (C₁-C₇) alkyl or A. The reaction is typically carried out at a temperature from about -78 to 40° C (preferably from about -78° C to room temperature) for about 0.5 to 48 hours (preferably for about 1.5 to 17 hours: 0.5 to 1 hour to generate the dianion of formula XI and 1 to 16 hours for the alkylation). The product of the reaction is an acid of the formula XII, wherein R² and R³ are as defined immediately above, and R⁴ is (C₁-C₇) alkyl or A. The acid of formula XII so formed may be converted to the corresponding amide of the formula ID, wherein R² and R³ are as defined immediately above and R⁴ is (C₁-C₇) alkyl or A, by the acid to amide synthesis illustrated in scheme 1 and described above.

40 Compounds of the formula ID, wherein R⁴ is XR¹⁰, are prepared by the same procedure as that described above for the preparation of compounds of the formula ID wherein R⁴ is (C₁-C₇) alkyl or A, with one modification. The dianion of formula XI is reacted with a compound of formula R¹⁰SSR¹⁰ instead of HalR⁴. This reaction produces an acid of the formula XII, wherein R⁴ is XR¹⁰. The acid of the formula XII so formed can then be converted to the corresponding amide of formula ID by the acid to amide synthesis illustrated in scheme 1 and described above.

55 The aminopyrimidine and aminopyridine intermediates used in the present invention are known in the literature or may be prepared by methods known in the art from intermediates that are known in the literature or commercially available. References for the preparation of many of the pyrimidine and pyridine intermediates can be found in the monographs "The Pyrimidines", ed. by D.J. Brown (1962) and "Pyridine and its Derivatives", ed. by R.A. Abramovitch (1961), Interscience Publishers, Inc., New York, N.Y., and their

supplements. The preparation of certain of these intermediates is described in greater detail below.

2,6-Disubstituted-5-amino-pyrimidine derivatives may be prepared by reacting the appropriately substituted 4,6-dihydroxypyrimidine with a nitrating agent such as fuming nitric acid in acetic acid at a temperature from about 15°C to about 40°C for a period of about 1 to about 5 hours. The resulting 5-nitropyrimidine is converted to the 2,4-dichloro-5-nitropyrimidine intermediate using a chlorinating agent such as phosphoryl chloride, alone or in the presence of a base, preferably diethylaniline, at a temperature from about 100 to about 115°C for a period of about 0.5 to about 2 hours. Procedures for carrying out these transformations are described in J. Chem. Soc., 3832 (1954).

The 2,6-bis(alkylthio)-5-nitropyrimidine derivatives may be prepared by reacting the appropriate dichloro intermediate with two equivalents of sodium alkylthiolate in a solvent such as dimethylformamide or, preferably, methanol, for about 4 to about 16 hours at a temperature from about 0 to about 30°C, preferably at ambient temperature. Monosubstitution of the dichloro intermediate is then accomplished by using one equivalent of nucleophile, at a reaction temperature of about 0 to about 100°C, depending on the reactivity of the nucleophile, in an inert solvent such as dimethyl formamide or tetrahydrofuran, for a period of about 4 to about 16 hours.

The resulting monochloro derivative is then reacted with one equivalent of a different nucleophile to yield a disubstituted derivative with different substituents on the carbon atoms at positions 2 and 4. The 2,6-disubstituted-5-nitropyrimidine is reduced using a reducing agent such as stannous chloride in concentrated hydrochloric acid or hydrogen gas with an appropriate catalyst, to yield the corresponding 5-aminopyrimidine derivative.

The novel pyridines of formula XXVIII and other 2,4-disubstituted-3-aminopyridine derivatives may be prepared by reacting the appropriate 2,4-dihydroxypyridine with a nitrating agent such as concentrated nitric acid at 80-100°C for 15-60 minutes. For example the preparation of 2,4-dihydroxy-6-methyl-3-nitropyridine is described in J. Heterocyclic Chem., 1970, 7, 389. The resulting 2,4-dihydroxy-3-nitropyridine is sequentially converted to the 2,4-dichloro-3-nitropyridine, 2,4-disubstituted-3-nitropyridine and 2,4-disubstituted-3-aminopyridine derivatives, using reaction conditions similar to those described above for the pyrimidine series.

The preparation of certain 5-aminoquinolines and 5-aminoisoquinolines used as reactants in scheme 1 is illustrated in scheme 5. Referring to scheme 5, 5-aminoquinolines and isoquinolines of the formulae XV and XVII may be prepared as follows. A quinoline or isoquinoline of the formula XIII is nitrated at the 5 position, respectively, by reacting it with a nitrating agent such as nitric acid or potassium nitrate with or without an acid catalyst such as sulfuric acid, for from about 2 to 16 hours at a temperature from about 0-100°C. The nitro compound of formula XIV so formed is then reduced using a reducing agent such as stannous chloride, iron, zinc, or hydrogen gas with an appropriate catalyst, with or without an acid catalyst such as hydrochloric acid, for from about 2 to 16 hours at a temperature from about 0-100°C, to yield the corresponding 5-aminoquinoline or 5-aminoisoquinoline of formula XV.

Compounds of the formula XVII, wherein R⁵ is -SR¹⁴ and is attached to the quinoline or isoquinoline ring at the 6 position, and wherein R¹⁴ is (C₁-C₆) alkyl, (C₅-C₇) cycloalkyl, phenyl (C₁-C₄) alkyl, phenyl, substituted phenyl, heteroaryl, or substituted heteroaryl, may be prepared as follows. A compound of the formula XIV, wherein R⁵ is -Cl and is attached to the quinoline or the isoquinoline ring at the 6 position, is reacted with a compound of the formula R¹⁴SH, wherein R¹⁴ is as defined above, and a base such as sodium hydride, or such compound of the formula XIV is reacted with a compound of the formula R¹⁴SNa, wherein R¹⁴ is as defined above, in an inert solvent such as tetrahydrofuran, for about 4 to 16 hours at a temperature of from about -10°C to room temperature. The preferred temperature is -10°C. This reaction yields a compound of the formula XVI, which is then converted to the corresponding 5-aminoquinoline or isoquinoline of the formula XVII by the method described above for reduction of compounds of formula XIV.

The urea compounds of the formula I, wherein Q is -NR¹⁷R¹⁸, may be prepared by reacting a secondary amine of the formula NHR¹⁷R¹⁸ with a compound of the formula R¹NCO. The reaction is typically carried out at ambient temperature in a hydrocarbon solvent such as hexane. Secondary amines of the formula NHR¹⁷R¹⁸ may be prepared by a variety of methods well known in the art. (See, e.g., Vogel's Textbook of Practical Organic Chemistry, Longmen, Inc., New York, pp. 564-575 (4 ed. 1978).

Except where otherwise noted, pressure is not critical in any of the above reactions. Preferred temperatures for the above reactions were stated where known. In general, the preferred temperature for each reaction is the lowest temperature at which product will be formed. The preferred temperature for a particular reaction may be determined by monitoring the reaction using thin layer chromatography.

The novel compounds of formula I and the pharmaceutically acceptable salts thereof are useful as inhibitors of acyl coenzyme A: cholesterol acyltransferase (ACAT). As such they inhibit intestinal absorption of cholesterol in mammals and are useful in the treatment of high serum cholesterol in mammals, including

humans. As used herein, treatment is meant to include both the prevention and alleviation of high serum cholesterol. The compound may be administered to a subject in need of treatment by a variety of conventional routes of administration, including orally, parenterally and topically. In general, these compounds will be administered orally or parenterally at dosages between about 0.5 and about 30 mg/kg body weight of the subject to be treated per day, preferably from about .08 to 5 mg/kg. For an adult human of approximately 70 kg of body weight, the usual dosage would, therefore, be about 3.5 to about 2000 mg per day. However, some variation in dosage will necessarily occur depending on the condition of the subject being treated and the activity of the compound being employed. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

A compound of formula I or a pharmaceutically acceptable salt thereof may be administered alone or in combination with pharmaceutically acceptable carriers, in either single or multiple doses. Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents. The resulting pharmaceutical compositions are then readily administered in a variety of dosage forms such as tablets, powders, lozenges, syrups, injectable solutions and the like. These pharmaceutical compositions can, if desired, contain additional ingredients such as flavorings, binders, excipients and the like. Thus, for purposes of oral administration, tablets containing various excipients such as sodium citrate, calcium carbonate and calcium phosphate may be employed along with various disintegrants such as starch, alginic acid and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in soft and hard filled gelatin capsules. Preferred materials for this include lactose or milk sugar and high molecular weight polyethylene glycols. When aqueous suspensions or elixirs are desired for oral administration, the essential active ingredient therein may be combined with various sweetening or flavoring agents, coloring matter or dyes and, if desired, emulsifying or suspending agents, together with diluents such as water, ethanol, propylene glycol, glycerin and combinations thereof.

For parenteral administration, solutions of a compound of formula I or a pharmaceutically acceptable salt thereof in sesame or peanut oil, aqueous propylene glycol, or in sterile aqueous solution may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. Such solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, the sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

The activity of the compounds of the present invention as ACAT inhibitors may be determined by a number of standard biological or pharmacological tests. For example, the following procedure was used to determine the ACAT inhibiting activity of compounds of formula I. ACAT was assayed in microsomes isolated from chow fed Sprague-Dawley rats according to Bilheimer, J.T., *Meth. Enzymol.*, 111, ps 286-293 (1985), with minor modifications. Microsomes from rat liver were prepared by differential centrifugation and washed with assay buffer prior to use. The assay mixture contained 25 μ l of BSA (40 mg/ml), 30 μ l of rat liver microsome solution (100 μ g microsomal protein), 20 μ l of assay buffer (0.1 M K_2PO_4 , 1.0 mM reduced Glutathione, pH 7.4), 20 μ g of cholesterol in 100 μ l of a 0.6% Triton WR-1339 solution in assay buffer, and 5 μ l of test compound dissolved in 100% DMSO (total volume = 180 μ l). The assay mixture was incubated for 30 min at 37°C. The reaction was started by the addition of 20 μ l of ^{14}C -Oleoyl-CoA (1000 μ M, 2,000 dpm/nmol) and run for 15 min at 37°C. The reaction was stopped by the addition of 1 ml ETOH. The lipids were extracted into 4 ml hexane. A 3 ml aliquot was dried under N_2 , and resuspended in 100 μ l of chloroform. 50 μ l of chloroform were spotted on a heat activated TLC plate and developed in hexane: diethyl ether: acetic acid (85:15:1, v:v:v). Incorporation of radioactivity into cholesteryl esters was quantified on a Berthold LB2842 Linear TLC Analyzer. ACAT inhibition was calculated relative to a DMSO control assay.

The activity of the compounds of formula I in inhibiting intestinal absorption of cholesterol may be determined by the procedure of Melchoir and Harwell, *J. Lipid. Res.*, 26, 306-315 (1985).

The present invention is illustrated by the following examples. It will be understood, however, that the invention is not limited to the specific details of these examples. Melting points are uncorrected. Proton nuclear magnetic resonance spectra (1H NMR) and C^{13} nuclear magnetic resonance spectra (C^{13} NMR) were measured for solutions in deuteriochloroform ($CDCl_3$) and peak positions are expressed in parts per million (ppm) downfield from tetramethylsilane (TMS). The peak shapes are denoted as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad; c, complex.

EXAMPLE 1

Ethyl 2-(4-n-Propylphenylthio)nonanoate

5

1.6 g (0.033 mole) sodium hydride (50% dispersion in mineral oil) was added to a solution of 5.0 g (0.033 mole) 4-propylthiophenol in 25 ml anhydrous dimethylformamide. After 15 minutes, 8.8 g (0.033 mole) ethyl 2-bromononanoate (prepared according to J. Labelled Compounds Radiopharm. 14, 713 (1978)) was added and the resulting mixture was stirred at room temperature overnight. The reaction mixture was then diluted with 150 ml ethyl acetate and the resulting mixture was washed with 5 x 60 ml water and then with 60 ml saturated aqueous sodium chloride solution. The ethyl acetate solution was dried over anhydrous sodium sulfate, filtered and concentrated in vacuo. The resulting oil was chromatographed on 600 g silica gel, eluting with 7:3 hexane/methylene chloride to yield 9.0 g (81% yield) of the desired product as an oil.

¹H NMR(CDCl₃): δ 0.88 (c,6H); 1.1-1.5 (c,total 12H) including 1.12 (t,3H); 1.54-1.93 (c,4H); 2.54 (t,2H); 3.56 (q,1H); 4.07 (q,2H); 7.1 (d,2H); 7.36 (d,2H).

EXAMPLE 1A

20

2-Hexylthiodecanoic Acid

17.3 g (0.36 mol) sodium hydride (50% dispersion in mineral oil) was added portionwise with stirring (gas evolution) to a solution of 26.8 ml. (0.19 mol) hexanethiol in 500 ml. anhydrous dimethylformamide. The mixture was stirred at room temperature for 30 min., then 45.2g (0.18 mol) 2-bromodecanoic acid was added dropwise with stirring, keeping the temperature of the reaction mixture below 45°C. The reaction mixture was stirred at room temperature under nitrogen overnight. The mixture was then diluted with 500 ml. water and the pH of the resulting mixture was adjusted to 1.5 with 6N aqueous hydrochloric acid solution. This mixture was extracted with 3 x 400 ml. ethyl acetate and the combined ethyl acetate extracts were washed with 5 x 700 ml. water and 1 x 500 ml. brine. The ethyl acetate solution was dried over anhydrous sodium sulfate, filtered and concentrated in vacuo. The resulting oil was chromatographed on 2 kg. silica gel, eluting with methylene chloride to yield 35g (67% yield) of the desired product as an oil.

35

EXAMPLE 1B

40

Resolution of 2-hexylthiodecanoic acid

2-Hexylthiodecanoyl chloride was prepared by the procedure of Example 4A. A solution of 2-hexylthiodecanoyl chloride (2.39 g., 7.8 mmol) in 20 ml methylene chloride was added slowly with stirring under nitrogen to a solution of (R)-(-)-2-phenylglycinol (1.08 g, 7.9 mmol) and 4-dimethylaminopyridine (0.96 g, 7.9 mmol) in 80 ml methylene chloride at 5°C. The reaction mixture was stirred at room temperature overnight. Methylene chloride (100 ml.) was then added and the resulting solution was washed sequentially with 100 ml 1N aqueous hydrochloric acid solution, 100 ml water, 100 ml saturated aqueous sodium bicarbonate solution and 100 ml brine. The organic phase was dried over anhydrous sodium sulfate and concentrated in vacuo to a solid residue (3.1 g). The diastereomers were separated by column chromatography on 800g silica gel using 1:1 hexane-diethyl ether as eluant. The less polar diastereomer (1.09 g, [α]_D^{RT} = -9.85° (CH₃OH); mp 98-100°C) and 0.99 g of the more polar diastereomer ([α]_D^{RT} = -9.46° (CH₃OH); mp 105-108°C) were obtained along with 0.36 g of a mixture of diastereomers (total yield 76%). A solution of the less polar diastereomer (900 mg, 2.2 mmol) in 42 ml 1,4-dioxane and 42 ml 6N aqueous sulfuric acid solution was heated at 105°C under nitrogen for 15 hours. The reaction mixture was cooled to room temperature, diluted with 80 ml water and the resulting mixture was extracted with 4 x 60 ml ethyl acetate. The combined ethyl acetate extracts were washed with 60 ml brine, dried over anhydrous sodium sulfate and concentrated in vacuo to yield (S)-(-)-2-hexylthiodecanoic acid as an oil (634 mg., 99.6% yield); [α]_D^{RT} =

55

-59.5° (CH₃OH).

In a similar manner, hydrolysis of the more polar diastereomer yielded 98.4% of (R)-(+)-2-hexylthiodecanoic acid as an oil; $[\alpha]_D^{25} = +54.0^\circ$ (CH₃OH).

5

EXAMPLE 2

10 Ethyl 2-(4-t-Butylphenylthio)octanoate

A mixture of 5.0 g (0.02 mole) ethyl 2-bromooctanoate, 3.37 g (0.02 mole) p-t-butylthiophenol and 3.31 g (0.24 mole) potassium carbonate in 70 ml acetone was refluxed under nitrogen overnight. The reaction mixture was cooled to room temperature and filtered and the filtrate was concentrated in vacuo. The residue was chromatographed on 500 g silica gel, eluting with 6:4 methylene chloride/hexane to yield 3.8 g (57% yield) of the desired product as an oil.

15

¹H NMR(CDCl₃): δ 0.88 (c,3H); 1.1-1.52 (c,total 20H) including 1.14 (t,3H) and 1.3 (s); 1.66-2.11 (c,2H); 3.58 (q,1H); 4.1 (q,2H); 7.36 (m,4H).

20

EXAMPLE 3

25 2-(4-n-Propylphenylthio)nonanoic acid

A solution containing 5.7 (0.017 mole) of the title compound of Example 1, 35 ml of 1N aqueous sodium hydroxide solution (0.035 mole) and 3 ml methanol was refluxed overnight. The resulting solution was cooled to room temperature, acidified to pH 1.5 with 2N aqueous hydrochloric acid and extracted with 3 x 50 ml ethyl acetate. The combined ethyl acetate extracts were washed with 50 ml water and 50 ml saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate and concentrated in vacuo to yield the title compound as an oil (5.0 g, 96% yield) which was used in the subsequent reaction without further purification.

30

¹H NMR(CDCl₂): δ 0.88 (c, 6H); 1.17-1.54 (c, 12H); 1.54-1.92 (c, 4H); 2.53 (t, 2H); 3.54 (t, 1H); 7.1 (d, 2H); 7.37 (d, 2H).

35

EXAMPLE 4

40

2-(4-n-Propylphenylthio)-N-(2,4,6-trimethoxyphenyl)nonanamide

1.54 g (5 mmole) of the title compound of Example 3 in 20 ml of thionyl chloride was refluxed for 3 hours and then concentrated to dryness in vacuo. 523 mg (1.6 mmole) of the resulting acid chloride was dissolved in 20 ml methylene chloride and to the solution was added 292 mg (1.6 mmole) 2,4,6-trimethoxyaniline and 195 mg (1.6 mmole) 4-dimethylaminopyridine. The resulting solution was stirred at room temperature overnight and then concentrated in vacuo. The residue was partitioned between 60 ml ethyl acetate and 20 ml 1N aqueous hydrochloric acid solution. The ethyl acetate layer was washed with 50 ml water and 50 ml saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate and concentrated in vacuo. The crude product was chromatographed on 100 g silica gel, eluting with 1:1 hexane/ethyl acetate to yield 370 mg (49% yield) of the title compound as a whitish solid.

50

55

EXAMPLE 4A

N-[2-methyl-4,6-bis(methylthio) pyrimidin-5-yl]-2-hexylthiodecanoic amide

A solution of 6.49 g (22.5 mmol) 2-hexylthiodecanoic acid in 40 ml thionyl chloride and 100 ml benzene was refluxed under nitrogen for 2.5 hours and then concentrated to dryness in vacuo. The resulting acid chloride (6.88 g, 22.5 mmol) was dissolved in 15 ml. methylene chloride and the solution was added dropwise to a solution of 4.63 g (23 mmol) 5-amino-4,6-bis(methylthio)-2-methylpyrimidine in 140 ml methylene chloride. The resulting solution was refluxed under nitrogen overnight. The reaction solution was then cooled, diluted with 140 ml methylene chloride and washed with 2 x 125 ml 3N aqueous hydrochloric acid solution, 1 x 125 ml water, 1 x 125 ml saturated aqueous sodium bicarbonate solution and 1 x 125 ml brine. The methylene chloride solution was dried over anhydrous sodium sulfate, filtered and concentrated to dryness in vacuo. The solid residue was recrystallized from diethyl ether yielding 5.35 g of the title compound, m.p. 99-101 °C. The filtrate was concentrated in vacuo and the residue was chromatographed on 400 g silica gel eluting with 9:1 hexane/ethyl acetate. Recrystallization of the product obtained by chromatography from diethyl ether yielded another 2.32 g of the title compound, m.p. 99 °-101 °C (total yield 72.4%).

¹H NMR (CDCl₃): δ 0.87 (c, 6H); 1.21-1.84 (c, 21 H), 2.02 (m, 1H); 2.50 (s, 6H); 2.76 (s, 3H), 2.74 (t, 2H); 3.45 (t, 1H), 8.08 (s, 1H);

IR (CHCl₃): 2923, 2852, 1681, 1511, 1468, 1431, 1405 cm⁻¹.

EXAMPLE 4BN-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-hexylthiodecanoic amide

A solution of 4.19 g (13.7 mmol) 2-hexylthiodecanoyl chloride, prepared according to Example 4A, in 15 ml methylene chloride was added dropwise with stirring under nitrogen to a solution of 2.75 g (13.7 mmol) 3-amino-2,4-bis(methylthio)-6-methylpyridine in 30 ml pyridine cooled to 5 °C. The reaction mixture was stirred at room temperature under nitrogen overnight. Methylene chloride (250 ml) was then added to the reaction mixture and the resulting solution was washed with 3 x 50 ml 3N aqueous hydrochloric acid solution, 2 x 50 ml water, 1 x 50 ml saturated aqueous sodium bicarbonate solution and 1 x 50 ml brine. The methylene chloride solution was dried over anhydrous sodium sulfate, filtered and concentrated to dryness in vacuo. The solid residue (6.5 g) was recrystallized from petroleum ether to yield 4.7 g of the title compound, m.p. 75-76.5 °C (72.8% yield).

¹H NMR (CDCl₃): δ 0.86 (c, 6H); 1.16-1.74 (c, 21H); 2.04 (m, 1H); 2.4 (s, 3H) 2.48 (s, 3H); 2.5 (s, 3H); 2.77 (t, 2H); 3.45 (t, 1H); 6.65 (s, 1H); 8.14 (s, 1H).

IR (CHCl₃): 2922, 2852, 1676, 1600, 1561, 1466 cm⁻¹.

EXAMPLE 52-Bromo-N-(2,4,6-trimethoxyphenyl)decanamide

2-Bromodecanoic acid (1 g, 3.8 mmol) was heated under reflux in thionyl chloride (10 ml) for 1 hour. The thionyl chloride was evaporated and the residue was dissolved in dry ether (10 ml) and added dropwise to a solution of 2,4,6-trimethoxyaniline (0.7 g, 3.8 mmol) in pyridine (20 ml) at 0 °C and the mixture was stirred for 1.5 hours. The reaction mixture was poured into saturated aqueous ammonium chloride and extracted three times with ethyl acetate (60 ml). The combined organics were extracted with water and brine and dried and concentrated. Recrystallization from isopropyl ether afforded 1.1g (65%) of the title compound, m.p. 109-110 °C. This material was used directly in the next step.

EXAMPLE 6

N-(2,4,6-Trimethoxy)phenyl-2-((2-pyridyl)thio)decanoamide

2-Thiopyridine (0.27 g, 2.4 mmol) in dimethylformamide (20 ml) was treated with sodium hydride (0.1 g, 2.4 mmol, 60% oil dispersion) and stirred for 15 minutes at 25 °C. To this cloudy solution, the title compound of Example 5 (1.0 g, 2.4 mmol) in dimethylformamide (10 ml) was added and the mixture and was stirred at 25 °C for 1.5 hours. The reaction mixture was then poured into 1 N HCl (75 ml) and extracted 3 times with ethyl acetate (125 ml). The organics were dried, concentrated and chromatographed on silica gel (eluted with 1:1 ethyl acetate:hexanes). Recrystallization from isopropyl ethyl afforded 0.4 g (36%) of the title compound, m.p. 92 °C.

¹H NMR (CDCl₃): δ 8.52-8.42 (m, 2H), 7.54 (t, J=4 Hz, 1H), 7.26 (m, 1H), 7.04 (t, J=4 Hz, 1H), 6.12 (s, 2H), 4.53 (t, J=3 Hz, 1H), 3.80 (s, 3H), 3.66 (s, 6H), 2.26-0.86 (m, 17H). IR (CHCl₃): 2920, 1685, 1595 cm⁻¹. Anal. Calculated for C₂₄H₃₄O₄N₂S: C, 64.55; H, 7.67; N, 6.27. Found: C, 64.34; H, 7.54; N, 6.20.

The title compounds of Examples 7 through 12 were prepared by a procedure similar to that described in Example 4.

EXAMPLE 72-(4-t-Butylphenylthio)-N-(2,4,6-trimethoxyphenyl)nonanamide (57% yield)

¹H NMR: δ 0.87 (c, 3H); 1.28 (s, 9H); 1.3 (c, 8H); 1.57 (c, 2H); 2.0 (c, 2H); 3.70 (s, 6H); 3.78 (s+c, 4H); 6.11 (s, 2H); 7.29 (d, 2H); 7.42 (d, 2H); 7.87 (s, 1H). IR (CHCl₃): 1672 cm⁻¹.

EXAMPLE 82-[4-(1,1-Dimethylpropyl)phenylthio]-N-(2,4,6-trimethoxyphenyl)nonanamide (60% yield)

¹H NMR: δ 0.65 (m, 3H); 0.87 (c, 3H); 1.24 (s, 6H); 1.28 (c, 8H); 1.62 (c, 4H); 1.97 (c, 2H); 3.71 (s, 6H); 3.75 (m, 1H); 3.78 (s, 3H); 6.12 (s, 2H); 7.22 (d, 2H); 7.41 (d, 2H); 7.89 (s, 1H). IR(CHCl₃): 1670 cm⁻¹.

EXAMPLE 92-(4-n-Butylphenylthio)-N-(2,4,6-trimethoxyphenyl)nonanamide (22% yield)

¹H NMR: δ 0.89 (m, 6H); 1.32 (c, 10H); 1.58 (c, 4H); 1.97 (c, 2H); 2.55 (m, 2H); 3.71 (s, 6H); 3.75 (m, 1H); 3.78 (s, 3H); 6.11 (s, 2H); 7.08 (d, 2H); 7.41 (d, 2H); 7.86 (s, 1H). IR (CHCl₃): 1670 cm⁻¹.

EXAMPLE 102-[4-(1-Methylpropyl)phenoxy]-N-(2,4,6-trimethoxyphenyl)nonanamide (50% yield)

¹H NMR: δ 0.87 (c, 6H); 1.2 (d, 3H); 1.3 (c, 8H); 1.55 (m, 4H); 2.0 (m, 2H); 2.55 (m, 1H); 3.70 (s, 6H); 3.77 (s, 3H); 4.6 (t, 1H); 6.1 (s, 2H); 6.96 (d, 2H); 7.09 (d, 2H); 7.38 (s, 1H). IR (CHCl₃): 1680 cm⁻¹.

EXAMPLE 11

2-(4-n-Propylphenoxy)-N-(2,4,6-trimethoxyphenyl)decanamide (59% yield)

¹H NMR: δ 0.88 (c, 6H); 1.28 (c, 10H); 1.6 (c, 4H); 2.0 (m, 2H); 2.52 (t, 2H); 3.71 (s, 6H); 3.77 (s, 3H); 4.59 (t, 1H); 6.1 (s, 2H); 6.95 (d, 2H); 7.1 (d, 2H); 7.37 (s, 1H). IR (CHCl₃): 1683 cm⁻¹.

5

EXAMPLE 12

10

2-(4-n-Propylphenylthio)-N-(2,4,6-trimethoxyphenyl)nonanamide (49% yield)

¹H NMR: δ 0.89 (m, 6H); 1.3 (c, 8H); 1.59 (m, 4H); 1.95 (c, 2H); 2.53 (t, 2H); 3.71 (s, 6H); 3.75 (m, 1H); 3.78 (s, 3H); 6.11 (s, 2H); 7.09 (d, 2H); 7.41 (d, 2H); 7.86 (s, 1H). IR (CHCl₃): 1670 cm⁻¹.

15 The title compounds of Examples 13 through 24 were prepared by a procedure similar to that described in Examples 5 and 6.

EXAMPLE 13

20

N-(2,4,6-Trimethyl)phenyl-2-(1-decylthio)octanamide

25 M.p. 42° C. ¹H NMR: δ 8.06 (s, 1); 6.85 (s, 2); 3.39 (t, 3 Hz, 1); 2.61 (t, 4 Hz, 2); 2.23 (s, 3); 2.16 (s, 6); 1.58 (m, 4); 1.23 (bs, 22); 0.84 (t, 4 Hz, 6). IR (CHCl₃): 3340, 2930, 1675, 1500 cm⁻¹.

EXAMPLE 14

30

N-(2,4,6-Trimethyl)phenyl-2-(pyrid-2-ylthio)decanamide

35 M.p. 85-87° C. ¹H NMR: δ 8.81 (s, 1); 8.36 (d, 2 Hz, 1); 7.50 (t, 3 Hz, 1); 7.24 (d, 3 Hz, 1); 7.01 (t, 3 Hz, 1); 6.80 (s, 2); 4.48 (t, 3 Hz, 1); 2.21 (s, 3); 2.0 (s, 6); 1.85-0.80 (m, 17). IR (CHCl₃): 2930, 1685, 1585 cm⁻¹.

EXAMPLE 15

40

N-(2,4,6-Trimethyl)phenyl-2-((2-methylfuryl)thio)decanoamide

45 M.p. 64-65° C. ¹H NMR: δ 7.92 (bs, 1); 7.34 (s, 1); 6.88 (s, 2); 6.28 (m, 1); 6.21 (d, 1 Hz, 1); 3.88 (s, 2); 3.42 (t, 3 Hz, 1); 2.26 (s, 3); 2.18 (s, 6); 2.04 - 0.82 (m, 17). IR (CHCl₃): 2930, 1675, 1495 cm⁻¹.

EXAMPLE 16

50

N-(2,4,6-Trimethyl)phenyl-2-[(2(6-ethoxy-benzothiazolyl)thio] octanamide

55 M.p. 106-108° C. ¹H NMR: δ 7.90 (s, 1); 7.60 (s, 1); 6.86 (s, 2); 4.83 (t, 3 Hz, 1); 4.48 (t, 3 Hz, 1); 3.32 (q, 4 Hz, 2); 2.37 (s, 3); 2.12 (s, 6) 2.0 - 0.85 (m, 26). IR (CHCl₃): 2920, 1680, 1595 cm⁻¹.

EXAMPLE 175 N-(2,4,6-Trimethyl)phenyl-2-[4-(7-trifluoromethylquinolinyl)thio] decanamide

M.p. 195-196 °C. ¹H NMR: δ 8.77 (d, 3 Hz, 1); 8.42 (s, 1); 8.27 (d, 5 Hz, 1); 7.84 (s, 1); 7.77 (d, 5 Hz, 1); 8.54 (d, 3 Hz, 1); 6.80 (s, 2); 4.26 (t, 3 Hz, 1); 2.22 (s, 3); 1.91 (s, 6); 2.20 - 0.80 (m, 17). IR (CHCl₃): 29,20, 1680, 1495 cm⁻¹.

10

EXAMPLE 18

15

N-(2,4,6-Trimethyl)phenyl-2-((2-thiazolyl)thio) decanamide

M.p. 74-75 °C. ¹H NMR: δ 8.82 (bs, 1); 7.25 (d, 1 Hz, 1); 6.84 (s, 2); 4.41 (t, 3 Hz, 1); 2.24 (s, 3); 2.05 (s, 6); 1.96 - 0.84 (m, 17). IR (CHCl₃): 2920, 1850, 1685, 1490 cm⁻¹.

20

EXAMPLE 19

25

N-(2,4,6-Trimethyl)phenyl-2-((2-quinolinyl)thio) decanamide

M.p. 11-112 °C. ¹H NMR: δ 9.14 (bs, 1); 7.98 (d, 4 Hz, 1); 7.89 (d, 3 Hz, 1); 7.77 (d, 3 Hz, 1); 7.65 (t, 3 Hz, 1); 7.47 (t, 3 Hz, 1); 7.31 (d, 4 Hz, 1); 6.82 (s, 2); 4.82 (t, 3 Hz, 1) 2.38 - 0.85 (m, 28). IR (CHCl₃): 2920, 2850, 1680, 1590 cm⁻¹.

30

EXAMPLE 20

35

N-(2,4,6-Trimethoxy)phenyl-2-(1-hexylthio) octanamide

M.p. 56-58 °C. ¹H NMR: δ 7.79 (s, 1); 6.12 (s, 2); 3.78 (s, 3); 3.76 (s, 76 (s, 6); 3.44 (t, 4 Hz, 1); 2.66 (m, 2); 1.90-0.87 (m, 24). IR (CHCl₃): 2930, 1675, 1490 cm⁻¹.

40

EXAMPLE 21

45

N-(2,4,6-Trimethoxy)phenyl-2-(1-decylthio) octanamide

M.p. 54-56 °C. ¹H NMR: δ (CDCl₃) 7.81 (s, 1); 6.15 (s, 2); 3.81 (s, 3); 3.79 (s, 6); 3.47 (t, 4 Hz, 1) 2.69 (m, 4); 1.63-1.92 (m, 6); 1.60 (m, 20); 0.90 (m, 6). IR (CHCl₃): 2920, 1670, 1600, 1460 cm⁻¹.

50

EXAMPLE 22

55

N-(2,4,6-Trimethyl)phenyl-2-(iso-butylthio) octanamide

M.p. 58-60° C. ¹H NMR: δ 8.03 (s, 1); 6.85 (s, 2) 3.37 (t, 4 Hz, 1); 2.52 (m, 2); 2.17 (s, 6); 1.83-0.86 (m, 16). IR (CHCl₃): 2930, 1670, 1495 cm⁻¹.

EXAMPLE 23

N-(2,4,6-Trimethylphenyl)-2-[2-(3-propyloxy)pyridyl]thio]decanamide

M.p. 68-69° C. ¹H NMR: δ 8.74 (bs, 1); 8.02 (m, 1); 7.02 (d, 1 Hz, 2); 6.82 (s, 2); 4.56 (t, 3 Hz, 1); 4.03 (t, 3 Hz, 2); 2.25 (s, 3); 2.02 (s, 6); 1.94 - 0.82 (m, 22). IR (CHCl₃): 2910, 2840, 1680, 1490 cm⁻¹.

EXAMPLE 24

N-(Isoquinolin-5-yl)-2-((2-pyridyl)thio) decanamide

M.p. 81-83° C. ¹H NMR: δ 9.24 (bs, 1); 8.60 - 8.36 (m, 3); 7.78 - 7.11 (m, 7); 4.53 (t, 3 Hz, 1); (CHCl₃): 2940, 2860, 1700, 1590 cm⁻¹.

EXAMPLE 25

N-(2,4,6-Trimethoxyphenyl)-2-methyl-2-(4-(1-methylpropyl)phenoxy)nonanoic amide

By use of the procedures described in Examples 1 and 3, ethyl 2-bromononanoate and 4-(1-methylpropyl)phenol were coupled and the product saponified to give 2-(4-(1-methylpropyl)phenoxy)-nonanoic acid. This material (1.0 g) was then methylated at the 2-position according to the procedure of Pfeffer, et. al. (J. Org. Chem., 1972, 37, 451) to give 2-methyl-2-hexanethiodecanoic acid (0.928 g). This material (0.86 g) was converted to the corresponding acid chloride with oxalyl chloride and coupled with 2,4,6-trimethoxyaniline (0.49 g) according to the procedure of Adams and Ulrich (J. Am. Chem. Soc., 1920, 42, 599) to give the title compound (1.12 g).

Oil. ¹H NMR: δ 7.82 (s, 1H); 7.12 (d, 6 Hz, 2 H); 7.07 (d, 6 Hz, 2 H); 6.19 (s, 2 H); 3.85 (s, 3 H); 3.83 (s, 6 H); 2.61 (dt, 8 Hz, 1 H); 1.98 (m, 2 H); 1.68 - 1.20 (m, 12 H); 1.52 (s, 3 H); 1.26 (d, 8 Hz, 3 H); 0.93 (m, 3 H); 0.85 (t, 8 Hz, 3 H). ¹³C NMR: δ 173.38, 159.90, 156.48, 152.69, 142.12, 127.39, 121.24, 107.15, 91.02, 84.41, 55.83, 55.47, 40.94, 40.06, 31.86, 31.32, 29.85, 29.33, 23.46, 22.68, 21.90, 21.67, 14.11, 12.21. IR (CHCl₃) cm⁻¹: 3410, 2940, 2850, 1680, 1608. Mass spectrum m/e (relative intensity): M+ 485.42 (16), 336.28 (33), 308.28 (24), 275.30 (30), 209.04 (40), 183.14 (100). High resolution mass spectra: m/e 485.3134, calcd for C₂₉H₄₃NO₅: 485.3141. Anal.: Calc'd for C₂₉H₄₃NO₅: C, 71.72; H, 8.93; N, 2.88. Found: C, 71.28; H, 8.87; N, 2.74.

EXAMPLE 26

N-(Isoquinolin-5-yl)-2-(4-(1-methylpropyl)phenoxy)nonanoic amide

N-(isoquinolin-5-yl)-2-bromodecanoic amide, prepared according to the procedures described in Example 3 and 25, was coupled with 4-(1-methyl propyl)phenol according to the procedure described in Example 6 to give the title compound.

Oil. ¹H NMR: δ 9.23 (s, 1 H); 8.61 (s, 1 H); 8.40 (d, 6 Hz, 1 H); 8.25 (d, 6 Hz, 1 H); 7.80 (d, 9 Hz, 1H); 7.62 (dd, 6 and 9 Hz, 1 H); 7.19 (d, 8 Hz, 2 H); 7.00 (d, 8 Hz, 2 H); 6.80 (d, 9 Hz, 1 H); 4.78 (t, 6 Hz, 1 H);

2.58 (tq, 6 & 9 Hz, 1 H); 2.12 (m, 2 H); 1.60 (m, 4 H); 1.25 (m, 11 H); 0.86 (t, 9 Hz, 3 H); 0.85 (t, 8 Hz, 3 H); ^{13}C NMR: δ 170.91, 155.34, 154.48, 152.94, 143.20, 142.07, 130.83, 129.60, 129.02, 128.58, 128.01, 127.40, 125.17, 124.26, 115.25, 113.49, 79.49, 40.94, 33.36, 31.80, 31.30, 29.35, 29.16, 25.26, 22.67, 22.08, 14.13, 12.20.

5 IR (CHCl₃) cm^{-1} : 3670, 3404, 2951, 2924, 1690, 1608, 1591. Mass spectrum m/e (relative intensity): $M + 432.2$ (16), 403.2 (7), 284.2 (30), 255.2 (36), 171.1 (25), 144.1 (100). Anal.: Calc'd for C₂₈H₃₆N₂O₂: C, 77.74; H, 8.39; N, 6.48. Found: C, 76.12; H, 8.57; N, 5.03.

10 EXAMPLE 27

15 N-(isoquinolin-5-yl)-2-(4-propylphenoxy)decanoic amide

N-(isoquinolin-5-yl)-2-bromodecanoic amide, prepared according to the procedures described in Examples 3 and 25, was coupled with 4-propylphenol according to the procedure described in Example 6 to give the title compound.

Oil. ^1H NMR: δ 9.24 (s, 1 H); 8.63 (s, 1 H); 8.41 (d, 5 Hz, 1 H); 8.24 (d, 7 Hz, 1 H); 7.80 (d, 8 Hz, 1 H); 20 7.60 (dd, 6 & 6 Hz, 1 H); 7.21 (d, 5 Hz, 1 H); 7.15 (d, 6 Hz, 2 H); 6.99 (d, 6 Hz, 2 H); 4.76 (t, 5 Hz, 1 H); 2.55 (t, 6 Hz, 2 H); 2.10 (m, 2 H); 1.62 (m, 4 H); 1.45-1.18 (br. m, 10 H); 0.94 (t, 5 Hz, 3 H); 0.85 (t, 4 Hz, 3 H). ^{13}C NMR: δ 170.77, 155.29, 153.01, 143.39, 136.99, 130.80, 129.94, 129.70, 129.00, 127.26, 125.02, 123.96, 115.24, 113.27, 79.54, 37.12, 33.28, 31.82, 29.38, 29.33, 29.19, 25.17, 24.70, 22.64, 14.08, 13.73. IR (CHCl₃) cm^{-1} : 3397, 2922, 1691, 1590. Mass spectrum m/e (relative intensity): $M + 432.30$ (4), 298.16 (24), 269.20 25 (36), 171.06 (22), 144.06 (100). Anal.: Calc'd for C₂₈H₃₆N₂O₂: C, 77.74; H, 8.39; N, 6.48. Found: C, 77.63; H, 8.43; N, 6.22.

30 EXAMPLE 28

35 N-(isoquinolin-5-yl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea

N-(4-(3-methylbutyl)phenyl)methyl-N-heptylamine was prepared and coupled with commercially available 5-aminoisoquinoline according to the procedure of DeVries, et. al. (J. Med. Chem. , 29 , 1131 (1986)) to give the title compound.

Oil. ^1H NMR: δ 9.15 (s, 1 H); 8.24 (d, 6 Hz, 1 H); 8.05 (d, 8 Hz, 1 H); 7.65 (d, 8 Hz, 1 H); 7.50 (dd, 8 & 8 Hz, 1 H); 7.32 (d, 8 Hz, 2 H); 7.28 (d, 8 Hz, 2 H); 6.67 (s, 1 H); 6.62 (d, 6 Hz, 1 H); 4.64 (s, 2 H); 3.52 (t, 6 40 Hz, 2 H); 2.70 (t, 6 Hz, 2 H); 1.75 (m, 2 H); 1.67-1.20 (m, 11 H); 0.95 (d, 6 Hz, 6 H); 0.88 (t, 6 Hz, 3 H). ^{13}C NMR: δ 155.72, 153.01, 143.51, 142.73, 134.38, 133.42, 129.46, 129.35, 129.07, 127.40, 126.91, 123.20, 123.02, 113.44, 51.34, 49.14, 41.03, 33.51, 31.87, 29.18, 28.79, 27.79, 27.08, 22.66, 22.59, 14.14. IR (CHCl₃) cm^{-1} : 3414, 2921, 2854, 1662, 1591, 1507. Mass spectrum m/e (relative intensity): $M + 445.3$ (6), 304.0 (7), 274.2 (2), 190.1 (20), 170.0 (32), 161.2 (100). High resolution mass spectra: m/e 445.3076, calc'd for 45 C₂₉H₃₉N₃O: 445.3093. Anal.: Calc'd for C₂₉H₃₉N₃O: C, 78.12; H, 8.82; N, 9.43. Found: C, 75.42; H, 8.59; N, 8.85.

50 EXAMPLE 29

55 N-(isoquinolin-5-yl)-2-(methoxycarbonylmethyl)nonadecanoic amide

Commercially available N-(isoquinolin-5-yl)-2-(carboxymethyl)nonadecanoic amide was esterified with diazomethane in ether to give the title compound.

^1H NMR: δ 9.16 (s, 1H); 8.57 (s, 1 H); 8.48 (d, 6 Hz, 1 H); 8.08 (d, 6 Hz, 1 H); 7.70 (d, 8 Hz, 1 H); 7.66 (d, 8 Hz, 1 H); 7.50 (dd, 6 and 8 Hz, 1 H); 5.50 (m, 1 H); 5.32 (m, 1 H); 3.70 (s, 3 H); 3.18-2.80 (m, 2 H);

2.70-2.20 (m, 3 H); 1.95 (m, 2 H); 1.20 (m, 22 H); 0.85 (t, 6 Hz, 3 H); ^{13}C NMR: δ 175.77, 170.66, 152.56, 142.68, 134.59, 134.39, 131.97, 128.87, 127.84, 125.35, 124.78, 114.77, 114.60, 51.93, 43.32, 41.85, 38.03, 35.28, 32.62, 32.54, 31.90, 29.68, 29.51, 29.43, 29.34, 29.19, 22.67, 14.11.

IR (KBr) cm^{-1} : 3418, 2918, 2848, 1724, 1692, 1591. Mass spectrum m/e (relative intensity): $M + 466.7$ (1), 305.3 (2), 259.2 (11), 226.1 (76), 186.1 (88), 171.0 (32), 144.0 (100). Anal.: Calc'd for $\text{C}_{29}\text{H}_{42}\text{NO}_3$: C, 74.96; H, 9.23; N, 5.83. Found: C, 74.88; H, 9.27; N, 5.78.

EXAMPLE 30

10

N-(Isoquinolin-5-yl)-2-(decyl)cyclopentane carboxamide

15 2-(Decyl)cyclopentane carboxylic acid, prepared according to the procedure of Hoeftle, et. al. (US 4,715,175), was coupled with 5-aminoisoquinoline according to the procedure outlined in Example 47 to give the title compound.

^1H NMR: δ 9.22 (s, 1 H); 8.51 (d, 5 Hz, 1 H); 8.10 (d, 7 Hz, 1 H); 7.78 (d, 7 Hz, 1 H); 7.75 (s, 1 Hz, H); 7.59 (d, 7 Hz, 1 H); 7.52 (dd, 5 & 7 Hz, 1 H); 2.26 (m, 2 H); 1.75 (m, 8 H); 1.23 (m, 16 H); 0.86 (t, Hz, 3 H).
20 ^{13}C NMR: δ 176.32, 153.13, 143.25, 131.98, 129.83, 129.04, 127.304, 124.70, 124.49, 113.60, 55.70, 40.40, 36.23, 31.87, 30.18, 29.58, 29.49, 20.30, 25.93, 24.74, 22.66, 14.09.

IR (KBr) cm^{-1} : 3432, 2923, 2851, 1686, 1591, 1506. Mass spectrum m/e (relative intensity): $M + 381.28$ (62), 240.08 (34), 209.20 (94), 144.08 (100). Anal. Calc'd for $\text{C}_{25}\text{H}_{36}\text{N}_2\text{O}$: C, 78.90; H, 9.54; N, 7.36. Found: C, 78.53; H, 9.58; N, 7.27.

25

EXAMPLE 31

30

N-(3-Methylquinolin-5-yl)-2-(4-(1-methylpropyl)phenoxy)nonanoic amide

3-Methyl-4-chloro-5-nitroquinoline was hydrogenated using Pd/C to give 3-methyl-5-aminoquinoline. This material was coupled with 2-(4-(1-methylpropyl)phenoxy)nonanoic acid according to the procedure outlined in Example 25 to give the title compound.

Oil. ^1H NMR: δ 8.70 (s, 1 H); 8.48 (s, 1 H); 7.93 (d, 9 Hz, 1 H); 7.88 (d, 9 Hz, 1 H); 7.64 (dd, 9 and 9 Hz, 1 H); 7.52 (s, 1 H); 7.18 (d, 9 Hz, 2 H); 7.01 (d, 9 Hz, 2 H); 4.78 (dd, 6 & 8 Hz, 1 H); 2.61 (tq, 9 and 9 Hz, 1 H); 2.40 (s, 3 H); 2.11 (m, 2 H); 1.72-1.26 (m, 12 H); 1.24 (d, 7 Hz, 3 H); 0.90 (m, 3 H); 0.84 (t, 10 Hz, 3 H).
40 ^{13}C NMR: δ 170.85, 155.39, 152.24, 147.70, 141.84, 130.65, 128.50, 128.17, 128.07, 127.21, 122.40, 121.62, 115.08, 79.43, 40.84, 33.39, 31.75, 31.31, 31.24, 29.32, 29.13, 25.31, 22.62, 21.91, 18.87, 14.06, 12.20. Mass spectrum m/e (relative intensity): $M + 446.32$ (11), 297.22 (59), 269.22 (85), 185.05 (44), 158.08 (100). High resolution mass spectra: m/e 446.2938, calc'd for $\text{C}_{29}\text{H}_{38}\text{N}_2\text{O}_2$: 446.2928. Anal.: Calc'd for $\text{C}_{29}\text{H}_{38}\text{N}_2\text{O}_2$: C, 77.99; H, 8.58; N, 6.27. Found: C, 75.94; H, 8.40; N, 6.65.

45

EXAMPLE 32

N-(2-Methyl-6-fluoroquinolin-5-yl)-2-(hexylthio)decanoic amide

2-Methyl-5-amino-6-fluoroquinoline, prepared by reduction of the corresponding nitro compound according to Example 31, was coupled with 2-hexylthiodecanoic acid according to the procedure of Example 25 to give the title compound.

50 ^1H NMR: δ 8.52 (s, 1 H); 7.95 (d, 10 Hz, 1 H); 7.91 (m, 1H); 7.46 (dd, 10 & 12 Hz, 1 H); 7.27 (d, 10 Hz, 1 H); 3.50 (t, 8 Hz, 1 H); 2.70 (s, 3 H); 2.68 (t, 7 Hz, 1 H); 2.10 (m, 1 H); 1.82 (m, 1 H); 1.70-1.16 (m, 20 H); 0.82 (t, 7 Hz, 6 H). ^{13}C NMR: δ 172.10, 158.50, 131.60, 129.80, 122.72, 119.40, 119.20, 51.22, 33.07, 32.10, 31.82, 31.33, 29.36, 29.26, 29.19, 28.59, 27.55, 25.10, 22.64, 22.49, 14.07, 13.97. IR (KBr) cm^{-1} : 3243,

2928, 2862, 1656. Mass spectrum m/e (relative intensity): M + 448.34 (1), 243.20 (8), 231.14 (9), 218.10 (6), 176.14 (100). Anal.: Calc'd for C₂₆H₃₉FN₂OS: C, 69.91; H, 8.80; N, 6.27. Found: C, 69.44; H, 8.82; N, 5.45.

5

EXAMPLE 33N-(6-Methoxyquinolin-5-yl)-2-(hexylthio)decanoic amide

10

Commercially available 6-methoxyquinoline (13.80 g) was nitrated according to the procedure of Campbell, et. al. (J. Am. Chem. Soc., 1946, 68, 1559) to give 5-nitro-6-methoxyquinoline (17.51 g). This crude product was directly reduced according to the procedure of Jacobs, et. al. (J. Am. Chem. Soc., 1920, 42, 2278) to give 5-amino-6-methoxyquinoline (6.25 g). This material (0.45 g) was coupled with 2-hexanethiodecanoic acid (0.75 g, prepared according to the procedures described in Examples 1 and 3) using the procedure described in Example 25 to give the title compound (0.63 g).

15

M.p. 88-89° C. ¹H NMR: δ 8.80 (d, 3 Hz, 1H); 8.59 (s, 1 H); 8.08 (d, 8 Hz, 1 H); 8.02 (d, 7 Hz, 1 H); 7.51 (d, 8 Hz, 1 H); 7.36 (d, 3 & 7 Hz, 1 H); 3.55 (t, 6 Hz, 1 H); 2.73 (t, 6 Hz, 2 H); 2.14-1.21 (m, 22H); 0.89 (t, 6 Hz, 6 H). ¹³C NMR: δ 172.10, 151.50, 148.50, 143.90, 131.52, 129.93, 126.00, 121.33, 118.30, 115.76, 56.37, 51.35, 33.23, 32.02, 31.86, 31.43, 29.43, 29.25, 29.35, 28.71, 27.62, 22.67, 22.54, 14.02. IR (KBr) cm⁻¹: 3233, 2920, 2849, 1659, 1526, 1501. Mass spectrum m/e (relative intensity): M + 444.28 (4), 328.22 (9), 243.18 (14), 229.08 (14), 216.06 (14), 174.20 (100). Anal.: Calc'd for C₂₆H₄₀N₂O₂S: C, 70.23; H, 9.07; N, 6.30. Found: C, 70.05; H, 9.03; N, 6.23.

20

25

EXAMPLE 34N-(6-methylthioquinolin-5-yl)-2-(hexylthio)decanoic amide

30

Commercially available 6-chloroquinoline (33.3 g) was nitrated according to the procedure described in Example 33 to give 5-nitro-6-chloroquinoline (20.36 g). This material (15 g) was allowed to react with sodium methylthiolate according to the procedure of Massie (Iowa State Coll. J. Sci. 1946, 21, 41; CA 41:3044 g) to give 5-nitro-6-methylthioquinoline (13.61 g). This material (3.70 g) was reduced using iron (5.62 g) and hydrochloric acid (1.5 ml) in 50% aqueous ethanol (50 ml) to give 5-amino-6-methylthioquinoline (3.0 g). This material (3.0 g) was coupled with 2-hexanethiodecanoic acid (5.83 g, prepared according to the procedures described in Examples 1 and 3) using the procedure described in Example 25, to give the title compound (3.8 g).

35

M.p. 91-92° C. ¹H NMR: δ 8.85 (d, 3 Hz, 1 H); 8.62 (s, 1 H); 8.05 (d, 9 Hz, 1 H); 8.00 (d, 9 Hz, 1 H); 7.65 (d, 9 Hz, 1 H); 7.40 (dd, 3 & 9 Hz, 1 H); 3.55 (t, 8 Hz, 1 H); 2.80 (t, 8 Hz, 2 H); 2.50 (s, 3 H); 2.10-1.35 (m, 17 H); 0.91 (t, 9 Hz, 6 H). ¹³C NMR: δ 172.00, 149.84, 131.37, 129.61, 126.91, 121.76, 51.22, 33.16, 32.36, 31.91, 31.47, 29.47, 29.34, 29.30, 28.69, 27.82, 22.73, 22.59, 15.77, 14.17, 14.08. IR (CHCl₃) cm⁻¹: 3318, 2923, 2852, 1877, 1586, 1567. Mass spectrum m/e (relative intensity): M + 460.2 (2), 413.2 (6), 344.2 (23), 295.2 (13), 243.2 (16), 217.0 (70), 190.1 (100). Anal.: Calc'd for C₂₆H₄₀N₂OS₂: C, 67.78; H, 8.75; N, 6.08. Found: C, 68.27; H, 8.46; N, 5.85.

40

45

EXAMPLE 35

50

N-(Quinolin-5-yl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea

5-Aminoquinoline was converted to the title compound according to the procedure described in Example 28.

55

Oil. ¹H NMR: δ 8.80 (d, 4 Hz, 1 H); 7.82 (d, 9 Hz, 1 H); 7.65 (dd, 6 & 9 Hz, 1 H); 7.61 (d, 6 Hz, 1 H); 7.48 (d, 9 Hz, 1 H); 7.26 (d, 6 Hz, 2 H); 7.22 (d, 6 Hz, 2 H); 7.15 (dd, 4 & 9 Hz, 1 H); 6.66 (s, 1 H); 4.53 (s, 2 H);

3.55 (t, 9 Hz, 2 H); 2.65 (t, 9 Hz, 2 H); 1.70 (m, 2 H); 1.60 (m, 3 H); 1.32 (m, 8 H); 1.00 (d, 6 Hz, 6 H); 0.88 (t, 6 Hz, 3 H). ^{13}C NMR: δ 156.03, 150.00, 148.69, 143.30, 134.54, 134.20, 129.89, 129.35, 129.31, 128.94, 126.96, 125.91, 120.73, 120.46, 51.19, 48.95, 41.05, 33.51, 31.87, 29.18, 28.75, 27.77, 27.06, 22.66, 22.60, 14.14. IR (CHCl_3) cm^{-1} : 3416, 2913, 2855, 1665, 1596, 1509. Mass spectrum m/e (relative intensity): $M + 445.3$ (4), 304.2 (6), 274.3 (7), 190.1 (8), 161.2 (100). High resolution mass spectra: m/e 445.3104, calc'd for $\text{C}_{29}\text{H}_{39}\text{N}_3\text{O}$: 445.3093. Anal.: Calc'd for $\text{C}_{29}\text{H}_{39}\text{N}_3\text{O}$: C, 78.12; H, 8.82; N, 9.43. Found: C, 75.75; H, 8.55; N, 9.00.

EXAMPLE 36

N-(6-Methoxyquinolin-5-yl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea

5-Amino-6-methoxyquinoline, prepared as described in Example 33, was converted to the title compound according to the procedure described in Example 28.

M.p. 86-87°C. ^1H NMR: δ 8.73 (d, 3 Hz, 1 H); 8.15 (d, 6 Hz, 1 H); 7.94 (d, 8 Hz, 1 H); 7.39 (d, 8 Hz, 1 H); 7.30 (m, 3 Hz, H); 7.22 (d, 6 Hz, 2 H); 6.44 (s, 1 Hz, H); 4.62 (s, 2 Hz, H); 3.80 (s, 3 Hz, H); 3.43 (t, 9 Hz, 2 H); 2.62 (t, 9 Hz, 2 H); 1.76-1.21 (m, 13 Hz, H); 0.95 (d, 6 Hz, 6 H); 0.88 (t, 6 Hz, 3 H). ^{13}C NMR: δ 156.97, 149.90, 148.27, 143.74, 142.49, 134.83, 132.39, 128.78, 128.09, 127.18, 126.12, 121.11, 120.80, 115.49, 56.25, 50.72, 48.08, 40.91, 33.45, 31.82, 29.12, 28.61, 28.45, 27.70, 26.96, 22.58, 22.54, 14.09. IR (CHCl_3) cm^{-1} : 3395, 2953, 2924, 1651, 1504. Mass spectrum m/e (relative intensity): $M + 475.4$ (6), 334.3 (11), 200.1 (14), 173.1 (17), 161.2 (100). Anal.: Calc'd for $\text{C}_{36}\text{H}_{41}\text{N}_3\text{O}_2$: C, 75.75; H, 8.69; N, 8.83. Found: C, 75.58; H, 8.90; N, 8.68.

EXAMPLE 37

N-(7-Methoxyisoquinolin-8-yl)-2-(4-(1-methylpropyl)phenoxy)nonanoic amide

7-Methoxy-8-aminoquinoline, prepared by reduction of the corresponding nitro compound according to Example 31, was coupled with 2-(4-(1-methylpropyl)phenoxy)nonanoic acid according to the procedure of Example 25 to give the title compound.

Oil. ^1H NMR: δ 9.14 (s, 1 H); 8.39 (d, 5 Hz, 1 H); 8.12 (s, 1 H); 7.73 (d, 7 Hz, 1 H); 7.54 (d, 5 Hz, 1 H); 7.44 (d, 7 Hz, 1 H); 7.16 (d, 6 Hz, 2 H); 7.01 (d, 6 Hz, 2 H); 4.74 (t, 5 Hz, 1 H); 3.79 (s, 3 H); 2.61 (tq, 10 and 10 Hz, 1 H); 2.14 (m, 2 H); 1.58 (m, 4 H); 1.46-1.22 (br, m, 8 H); 1.20 (d, 5 Hz, 3 H); 0.86 (m, 6 H). ^{13}C NMR: δ 172.16, 155.80, 151.89, 148.40, 141.53, 140.95, 131.04, 128.19, 127.30, 125.60, 119.96, 118.70, 115.40, 115.20, 79.93, 56.38, 40.89, 33.66, 31.79, 31.27, 29.38, 29.19, 25.36, 22.64, 22.00, 14.09, 12.25. IR (CHCl_3) cm^{-1} : Mass spectrum m/e (relative intensity): $M + 462.30$ (6), 314.22 (100), 285.22 (50), 229.12 (59).

EXAMPLE 38

N-(2-Phenyl-4-methoxycarbonylquinolin-3-yl)-2-(4-(1-methylpropyl)phenoxy)nonanoic amide

2-Phenyl-4-methoxycarbonylquinolin-3-ylquinoline was coupled with 2-(4-(1-methylpropyl)phenoxy)nonanoic acid according to Example 25 to give the title compound.

M.p. 89-91°C. ^1H NMR: δ 8.56 (s, 1 H); 8.14 (d, 9 Hz, 1 H); 8.05 (d, Hz, 1 H); 7.71 (dd, 7 and 7 Hz, 1 H); 7.58 (dd, 7 & 7 Hz, 1 H); 7.52 (d, 7 Hz, 2 H); 7.36 (m, 3 H); 7.07 (d, 8 Hz, 2 H); 6.69 (d, 8 Hz, 2 H); 4.41 (t, 6 Hz, 1 H); 3.89 (s, 3 H); 2.55 (tq, 7 and 7 Hz, 1 H); 1.74 (m, 2 H); 1.58 (m, 2 H); 1.10-1.40 (br, m, 10 H); 0.84 (m, 6 H). ^{13}C NMR: δ 170.78, 166.20, 155.56, 146.30, 141.51, 138.40, 132.60, 129.90, 129.66, 128.96, 128.68, 128.61, 128.17, 127.95, 124.95, 123.81, 121.70, 115.18, 52.76, 40.87, 32.93, 31.73, 31.25, 29.04,

25.03, 22.61, 22.01, 14.08, 12.24. IR (CHCl₃) cm⁻¹: 3410, 2960, 1725, 1680, 1620. Mass spectrum m/e (relative intensity): M + 566.36 (13), 417.24 (29), 389.26 (23), 357.22 (13), 205.12 (45), 279.12 (100). Anal.: Calc'd for C₃₆H₄₂N₂O₄: C, 76.30; H, 7.47; N, 4.94. Found: C, 76.01; H, 7.55; N, 4.91.

5

EXAMPLE 39N-(3-Methoxypyridin-2-yl)-2-(4-(1-methylpropyl)phenoxy)nonanoic amide

3-Methoxy-2-aminopyridine, prepared by reduction of the corresponding nitro compound according to Example 31, was coupled with 2-(4-(1-methylpropyl)phenoxy)nonanoic acid according to Example 25 to give the title compound.

Oil. ¹H NMR: δ 8.90 (s, 1 H); 8.05 (d, 3 Hz, 1 H); 7.25 (m, 3 H); 6.97 (d, 3 Hz, 1 H); 6.89 (d, 9 Hz, 2 H); 4.64 (t, 7 Hz, 1 H); 3.74 (s, 3 H); 2.50 (tq, 12 & 12 Hz, 1 H); 1.98 (m, 2 H); 1.51 (m, 4 H); 1.18-1.08 (br, m, 11 H); 0.84 (t, 4 Hz, 3H); 0.76 (t, 5 Hz, 3 H). ¹³C NMR: δ 172.40, 156.00, 140.10, 128.15, 119.86, 117.34, 115.45, 55.68, 40.84, 33.20, 32.30, 31.75, 31.24, 29.34, 29.11, 27.60, 25.26, 22.61, 21.70, 14.06, 12.19. IR (CHCl₃) cm⁻¹: 3387, 2922, 2854, 1702, 15.98. Mass spectrum m/e (relative intensity): M + 412.34 (8), 313.22 (41), 263.22 (100), 151.08 (30). Anal.: Calc'd for C₂₅H₃₆N₂O₃: C, 72.78; H, 8.80; N, 6.80. Found: C, 71.49, H, 8.88; N, 6.03.

25

EXAMPLE 40N-(2-Methoxy-4-methylpyridin-2-yl)-2-(4-(1-methylpropyl)phenoxy)nonanoic amide

3-Nitro-4-methyl-2-pyridone was methylated with methyl iodide and reduced with zinc and acetic acid to give 2-methoxy-3-amino-4-methylpyridine. This material was coupled with 2-(4-(1-methylpropyl)phenoxy)nonanoic acid according to the procedure of Example 25 to give the title compound.

Oil. ¹H NMR: δ 8.19 (s, 1 H); 7.10 (d, 7 Hz, 2 H); 7.04 (d, 5 Hz, 1 H); 6.93 (d, 7 Hz, 2 H) 6.03 (d, 5 Hz, 1 H); 4.63 (t, 6 Hz, 1 H); 3.48 (s, 3 H); 2.53 (tq, 11 & 11 Hz, 1 H); 2.07 (s, 3 H); 2.03 (m, 2 H); 1.55 (m, 4 H); 1.28 (m, 8 H); 1.18 (d, 6 Hz, 3 H); 0.87 (m, 3 H); 0.79 (t, 5 Hz, 3 H). ¹³C NMR: δ 170.87, 159.58, 155.83, 143.55, 141.21, 133.70, 128.08, 124.35, 115.47, 109.06, 40.84, 37.44, 33.37, 31.75, 31.28, 31.25, 29.28, 29.07, 25.19, 22.61, 21.92, 19.47, 14.08, 12.20. IR (CHCl₃) cm⁻¹: 2920, 2852, 1685, 1655, 1606. Mass spectrum m/e (relative intensity): M + 426.32 (10), 327.16 (7), 277.20 (52), 249.20 (35), 165.18 (100). Anal.: Calc'd for C₂₆H₃₈N₂O₃: C, 73.20; H, 8.98; N, 6.57. Found: C, 73.06; H, 9.11; N, 6.28.

40

EXAMPLE 41

45

N-(2-Methoxy-4-methylpyridin-2-yl)-2-(hexylthio)decanoic amide

3-Nitro-4-methyl-2-pyridone was methylated with methyl iodide and reduced with zinc and acetic acid to give 2-methoxy-3-amino-4-methylpyridine. This material was coupled with 2-hexylthiodecanoic acid according to the procedure of Example 25 to give the title compound.

M.p. 83-85 °C. ¹H NMR: δ 8.55 (s, 1 H); 7.04 (d, 6 Hz, 1 H); 6.07 (d, 6 Hz, 1 H); 3.54 (s, 3 H) 3.41 (t, 6 Hz, 1 H); 2.12 (s, 3 H); 2.03-1.17 (br, m, 22 H); 0.84 (t, 5 Hz, 3 H). ¹³C NMR: δ 171.36, 159.74, 142.90, 133.40, 125.06, 109.20, 50.91, 37.47, 33.01, 31.82, 31.73, 31.38, 29.33, 29.27, 29.25, 28.52, 27.55 22.66, 22.52, 19.51, 14.10, 14.03. IR (KBr) cm⁻¹: 3232, 2920, 2850, 1652, 1592. Mass spectrum m/e (relative intensity): M + 408.38 (5), 292.30 (16), 193.12 (17), 165.10 (54), 138.22 (100). Anal.: Calc'd for C₂₃H₄₀N₂O₂S: C, 67.60; H, 9.87; N, 6.86. Found: C, 67.56; H, 9.56; N, 6.58.

EXAMPLE 425 N-(2-Methoxy-4-methylpyridin-2-yl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea

3-Nitro-4-methyl-2-pyridone was methylated with methyl iodide and reduced with zinc and acetic acid to give 2-methoxy-3-amino-4-methylpyridine. This material was converted to the title compound according to the procedure described in Example 28.

10 M.p. 90-91 °C. ¹H NMR: δ 7.20 (d, 15 Hz, 2 H); 7.16 (d, 15 Hz, 2 H); 6.95 (d, 6 Hz, 1 H); 6.82 (s, 1 H); 6.08 (d, 6 Hz, 1 H); 4.53 (s, 2 H); 3.53 (s, 3 H); 3.31 (t, 6 Hz, 2 H); 2.58 (t, 6 Hz, 2 H); 2.15 (s, 3 H); 1.70-1.43 (br, m, 8 H); 1.24 (m, 10 H); 0.90 (d, 5 Hz, 6 H); 0.86 (t, 4 Hz, 3 H). ¹³C NMR: δ 159.98, 155.75, 142.66, 140.27, 135.04, 131.35, 128.63, 128.33, 128.07, 127.54, 127.32, 109.91, 53.87, 50.39, 49.55, 47.44, 40.89, 40.81, 37.48, 33.43, 31.84, 31.78, 30.12, 29.25, 29.02, 28.17, 27.66, 27.35, 26.99, 22.63, 22.54, 19.48, 14.09.
15 IR (KBr) cm⁻¹: 2952, 2922, 1660, 1635, 1590. Mass spectrum m/e (relative intensity): M+ 439.40 (14), 298.28 (10), 274.30 (26), 190.20 (40), 165.08 (81), 161.14 (100). Anal.: Calc'd for C₂₇H₄₁N₃O₂: C, 73.76; H, 9.40; N, 9.56. Found: C, 73.85; H, 9.25; N, 9.35.

20 EXAMPLE 4325 N-(Imidazo[1,2-a]pyridin-3-yl)-2-(4-(1-methylpropyl)phenoxy)nonanoic amide

3-Aminoimidazo[1,2-a]pyridine, synthesized by reduction of the corresponding nitro compound according to Example 31, was coupled with 2-(4-(1-methylpropyl)phenoxy)nonanoic acid according to the procedure of Example 25 to give the title compound.

Oil. ¹H NMR: δ 8.78 (s, 1 H); 7.72 (d, 6 Hz, 1 H); 7.62 (d, 8 Hz, 1 H); 7.55 (s, 1 H); 7.24 (d, 9 Hz, 2 H);
30 7.20 (dd, 7 and 7 Hz, 1 H); 6.94 (d, 9 Hz, 2 H); 6.85 (dd, 7 and 7 Hz, 1 H); 4.52 (t, 6 Hz, 1 H); 2.51 (tq, 11 and 11 Hz, 1 H); 1.94 (m, 2 H); 1.50 (m, 4 H); 1.20-1.10 (br m, 11 H); 0.90-0.70 (br. m, 6 H). ¹³C NMR: δ 172.45, 156.14, 148.75, 144.99, 140.63, 128.32, 127.90, 127.84, 121.41, 121.17, 120.89, 117.20, 114.91, 76.93, 61.03, 41.15, 34.43, 32.94, 31.23, 29.14, 28.69, 25.01, 22.66, 21.93, 21.81, 14.16, 12.19. IR (KBr) cm⁻¹: Mass spectrum m/e (relative intensity): M+ 421.28 (25), 272.18 (51), 159.04 (58), 133.04 (100).
35 Anal.: Calc'd for C₂₆H₃₅N₃O₂: C, 74.25; H, 8.39; N, 9.99. Found: C, 73.82; H, 9.02; N, 9.56.

40 EXAMPLE 4445 N-(Imidazo[1,2-a]pyridin-3-yl)-2-(hexylthio) decanoic amide

3-Aminoimidazo[1,2-a]pyridine, synthesized by reduction of the corresponding nitro compound according to the procedure of Example 31, was coupled with 2-hexylthiodecanoic acid according to Example 25 to give the title compound.

¹H NMR (CDCl₃): δ 8.66 (s, 1H); 7.75 (d, 5Hz, 1H); 7.62 (d, 8Hz, 1H); 7.52 (s, 1H); 7.22 (dd, 7 & 8Hz, 1H); 6.84 (dd, 5 & 7Hz, 1H); 3.48 (t, 6Hz, 1H); 2.67 (t, 6Hz, 2H); 2.03 (m, 1H); 1.83 (m, 1H); 1.70 - 1.15 (m, 20H); 0.85 (m, 6H).

50 EXAMPLE 4555 N-(8-Chloro-6-methoxyquinolin-5-yl)-2-hexylthiodecanoic amide

5-Amino-6-methoxy-8-chloroquinoline, produced as a side product of the reduction procedure described

in Example 33, was coupled with 2-hexylthiodecanoic acid according to the procedure described in Example 25 to give the title compound.

Mp = 110-111 °C. Anal.: Found: C, 65.40; H, 8.06; N, 5.73. Calc'd for $C_{25}H_{39}ClN_2O_2S$: C, 65.18; H, 8.42; N, 5.85.

EXAMPLE 46

N-(6,8-Di(methylthio)quinolin-5-yl)-2-hexylthiodecanoic amide

5-Amino-6,8-di(methylthio)quinoline, produced as a side product of the procedure described in Example 34, was coupled with 2-hexylthiodecanoic acid according to the procedure described in Example 25 to give the title compound.

Mp = 91-93 °C.

EXAMPLE 47

N-(6-Methylthioquinolin-5-yl)-2-(4-sec-butylphenoxy)nonanoic amide

5-Amino-6-methylthioquinoline, prepared as described in Example 34, was coupled with 2-(4-sec-butylphenoxy)nonanoic acid according to the procedure described in Example 25 to give the title compound.

Oil. Anal.: Found: C, 71.35; H, 7.98; N, 5.54. Calc'd. for $C_{29}H_{38}N_2O_2S$: C, 72.76; H, 8.00; N, 5.85.

EXAMPLE 48

N-(6-Methylthioquinolin-5-yl)-2-octanyl-1,3-dithiane-2-carboxamide

5-Amino-6-methylthioquinoline, prepared as described in Example 34, was coupled with 2-octanyl-1,3-dithiane-2-carboxylic acid, prepared by treatment of 1,3-dithiane-2-carboxylic acid with sodium hexamethyldisilazide and octanyl bromide, according to Example 25 to give the title compound.

Oil. Anal.: Found: C, 59.11; H, 5.81; N, 6.07. Calc'd. for $C_{23}H_{32}N_2OS_3$: C, 61.57; H, 7.19; N, 6.24.

EXAMPLE 49

N-(6-ethoxyquinolin-5-yl)-2-hexylthiodecanoic amide

6-Hydroxyquinoline was treated with sodium hydride and ethyl iodide to give 6-ethoxyquinoline. This material was nitrated, reduced and coupled with 2-hexylthiodecanoic acid according to the procedure of Example 33 to give the title compound.

Mp = 88-90 °C. Anal.: Found: C, 70.37; H, 9.01; N, 6.26. Calc'd. for $C_{27}H_{42}N_2O_2S$: C, 70.69; H, 9.23; N, 6.11.

EXAMPLE 50

N-(6-fluoroquinolin-5-yl)-2-hexylthiodecanoic amide

6-Fluoroquinoline, prepared according to the procedure of Sveinbjornsson et. al. (J. Org. Chem., 1951, 16, 1450), was nitrated, reduced and coupled with 2-hexylthiodecanoic acid according to the procedure of Example 33 to give the title compound.

Mp = 74-75° C. Anal.: Found: C, 69.04; H, 8.55; N, 6.57. Calc'd. for C₂₅H₃₇FN₂OS: C, 69.40; H, 8.62; N, 6.48.

The title compounds of Examples 51-53 were prepared according to the procedure described in Example 4.

EXAMPLE 514,5-Dimethyl-trans-2-n-heptyl-N-(2,4,6-trimethoxyphenyl)-cyclohex-4-enecarboxamide

72% yield. IR(CHCl₃): 1675 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.62 (s) and 1.12-2.48 (c) (total 24H); 3.78 (s, 6H); 3.79 (s, 3H); 6.13 (s, 2H); 6.48 (s, 1H).

EXAMPLE 524,5-Dimethyl-trans-2-n-nonyl-N-(2,4,6-trimethoxyphenyl)cyclohex-4-enecarboxamide

66% yield. IR(CHCl₃): 1676 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.62 (s) and 1.12-2.48 (c) (total 28 H); 3.78 (s, 6H); 3.79 (s, 3H); 6.13 (s, 2H); 6.48 (s, 1H).

EXAMPLE 534,5-Dimethyl-trans-2-n-octyl-N-(2,4,6-trimethoxyphenyl)cyclohex-4-enecarboxamide

46% yield. IR(CHCl₃): 1676 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.62 (s) and 1.12-2.48 (c) (total 26 H); 3.78 (s, 6H); 3.79 (s, 3H); 6.13 (s, 2H); 6.48 (s, 1H).

Using the procedure described in Example 4, the amides in the bicyclo[2.2.1]hept-5-ene and bicyclo[2.2.2]oct-5-ene series of Examples 54-59 were obtained as mixtures of *endo* - and *exo* - isomers which could be separated by column chromatography on silica gel, eluting with hexane/ethyl acetate.

EXAMPLE 543-n-Nonyl-endo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.1]hept-5-ene-2-carboxamide

30% yield. IR(CHCl₃): 1663 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.15-1.62 (c, 18H); 1.79 (c, 1H); 2.51 (c, 1H); 2.61 (c, 1H); 3.16 (c, 1H); 3.77 (s, 6H); 3.78 (s, 3H); 6.11 (s) and 6.14 (c) (total 3H); 6.3 (c, 1H); 6.36 (s, 1H).

EXAMPLE 55

5 Endo-3-n-nonyl-exo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.1]hept-5-ene-2-carboxamide

26% yield. IR(CHCl₃): 1678 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.14-1.45 (c, 17H); 1.66 (c, 1H); 1.91 (c, 1H); 2.4 (c, 1H); 2.8 (c, 1H); 3.0 (c, 1H); 3.78 (s, 9H); 6.13 (s + c, 3H); 6.21 (c, 1H); 6.42 (s, 1H).

10

EXAMPLE 56

15

Exo-3-n-octyl-endo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.2]oct-5-ene-2-carboxamide

20% yield. IR(CHCl₃): 1666 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.02-1.87 (c, 19H); 2.11 (c, 1H); 2.45 (c, 1H); 2.84 (c, 1H); 3.76 (s, 6H); 3.78 (s, 3H); 20 6.11 (s, 2H); 6.3 (c, 1H); 6.42 (s, 1H); 6.5 (c, 1H).

EXAMPLE 57

25

Endo-3-n-octyl-exo-N-(2,4,6-trimethoxyphenyl)bicyclo[2.2.2]oct-5-ene-2-carboxamide

24% yield. IR(CHCl₃): 1680 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.06 (c, 1H); 1.25 (c, 14H); 1.65 (c, 2H); 1.89 (c, 2H); 2.16 (c, 1H); 2.46 (c, 1H); 2.78 (c, 1H); 3.77 (s, 6H); 3.79 (s, 3H) 6.13 (s, 2H); 6.21 (c, 1H); 6.32 (c, 1H); 6.43 (s, 1H).

EXAMPLE 58

35

Exo-2-n-nonyl-endo-N-(2,4,6-trimethoxyphenyl)bicyclo[2.2.1]hept-5-ene-2-carboxamide

40 40 % yield. IR(CHCl₃): 1677 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.27 (c, 12H); 1.4-1.7 (c, 6H); 1.88 (c, 1H); 2.02 (c, 1H); 2.82 (c, 2H); 3.76 (s, 6H); 3.77 (s, 3H); 6.12 (s, 2H); 6.2 (c, 2H); 6.4 (s, 1H).

EXAMPLE 59

45

Endo-2-n-nonyl-exo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.1]hept-5-ene-2-carboxamide

50

2 % yield. IR 1675 cm⁻¹. ¹H NMR: δ 0.84 (t, 3H); 1.12-2.06 (c, 19H); 2.45 (c, 1H); 2.82 (c, 1H); 3.17 (c, 1H); 3.77 (s, 6H); 3.79 (s, 3H); 6.05-6.27 (c, 4H); 6.6 (s, 1H).

The title compounds of Examples 60-64 were prepared according to the procedure described in Example 4.

55

EXAMPLE 60

2-n-Nonyl-N-(2,4,6-trimethoxyphenyl)indane-2-carboxamide

72 % yield. IR(CHCl₃): 1676 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.24 (c, 12H); 1.49 (c, 3H); 1.75 (c, 2H); 2.97 (d, 2H); 3.54 (d, 2H); 3.74 (s, 6H); 3.78
5 (s, 3H); 6.12 (s, 2H); 6.51 (s, 1H); 7.2 (c, 4H).

EXAMPLE 61

10

2-n-Octyl-N-(2,4,6-trifluorophenyl)-1,2,3,4-tetrahydro-2-naphthamide

22 % yield. ¹H NMR: δ 0.86 (t, 3H); 1.15-1.6 (c, 13H); 1.9 (c, 2H); 2.2 (c, 1H); 2.88 (c, 3H); 3.27 (d, 1H);
15 6.66 (c, 3H); 7.13 (s, 4H).

EXAMPLE 62

20

2-n-Octyl-N-(2,4,6-trimethoxyphenyl)-1,2,3,4-tetrahydro-2-naphthamide

65 % yield. ¹H NMR: δ 0.87 (t, 3H); 1.29 (c, 10H); 1.5 (c, 3H); 1.86 (c, 2H); 2.22 (c, 1H); 2.8 (c, 2H); 3.0
25 (c, 1H); 3.3 (d, 1H); 3.66 (s, 6H); 3.75 (s, 3H); 6.06 (s, 2H); 6.56 (s, 1H); 7.10 (s, 4H).

EXAMPLE 63

30

2-n-Decyl-N-(2,4,6-trimethoxyphenyl)indane-2-carboxamide

61 % yield. IR(CHCl₃) 1675 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.23 (c, 14H); 1.49 (c, 2H); 1.75 (c, 2H); 2.97 (d, 2H); 3.54 (d, 2H); 3.74 (s, 6H); 3.78
35 (s, 3H); 6.12 (s, 2H) 6.5 (s, 1H); 7.2 (c, 4H).

EXAMPLE 64

40

2-n-Decyl-N-(2,4,6-trifluorophenyl)indane-2-carboxamide

33 % yield. IR(CHCl₃) 1695 cm⁻¹.

¹H NMR: δ 0.86 (t, 3H); 1.23 (c, 14H); 1.42 (c, 2H); 1.78 (c, 2H); 3.02 (d, 2H); 3.5 (d, 2H); 6.72 (c, 3H); 7.22
45 (c, 4H).

50

EXAMPLE 652-n-Nonyl-N-(2,4,6-trifluorophenyl)-1,2,3,4-tetrahydro-2-naphthamide

55

To a solution of 1.0 g (6.8 mmole) 2,4,6-trifluoroaniline and 830 mg (6.8 mmole) 4-dimethylaminopyridine in 40 ml methylene chloride cooled to 5 °C under nitrogen was added a solution of 2.12 g (6.6 mmole) 2-n-nonyl-1,2,3,4-tetrahydro-2-naphthoyl chloride in 10 ml methylene chloride. The

resulting solution was stirred at room temperature for 44 hours. Fifty milliliters of methylene chloride was then added and the solution was washed sequentially with 30 ml aqueous hydrochloric acid, 30 ml water and 30 ml brine. The methylene chloride solution was dried over anhydrous sodium sulfate and concentrated to dryness in vacuo. The residue (2.5 g) was purified by column chromatography on silica gel eluting with 2:1 methylene chloride-hexane to yield 1.66 g (58% yield) of the desired product as a white, low melting solid.

IR(CHCl₃) 1691 cm⁻¹. ¹H NMR: δ 0.87 (t, 3H); 1.16-1.6 (c, 15H); 1.9 (c, 2H); 2.2 (c, 1H); 2.9 (c, 3H); 3.27 (d, 1H); 6.67 (m, 2H); 6.73 (s, 1H); 7.13 (s, 4H).

EXAMPLE 66

2-n-Nonyl-N-(2,4,6-trimethoxyphenyl)-1,2,3,4-tetrahydro-2-naphthamide

The title compound was prepared according to the procedure described in Example 65, except that 422 mg 2-n-nonyl-1,2,3,4-tetrahydro-2-naphthoyl chloride, 247 mg (1.3 mmole) 2,4,6-trimethoxyaniline, and 165 mg (1.3 mmole) 4-dimethylaminopyridine in 12 ml methylene chloride were stirred at room temperature for 20 hours. There was obtained 421 mg product (68% yield).

IR(CHCl₃) 1670 cm⁻¹. ¹H NMR: δ 0.87 (t, 3H); 1.25 (c, 12H); 1.5 (c, 3H); 1.85 (c, 2H); 2.21 (c, 1H); 2.8 (c, 2H); 3.0 (c, 1H); 3.3 (d, 1H); 3.66 (s, 6H); 3.75 (s, 3H); 6.06 (s, 2H); 6.56 (s, 1H); 7.1 (s, 4H).

EXAMPLE 67

2-n-Nonyl-N-(2,4,6-trifluorophenyl)indane-2-carboxamide

The title compound was prepared according to the procedure described in Example 65, except that 398 mg (1.3 mmole) 2-n-nonylindane-2-carbonyl chloride, 220 mg (1.5 mmole) 2,4,6-trifluoroaniline, and 187 mg (1.5 mmole) 4-dimethylaminopyridine in 12 ml methylene chloride were stirred at room temperature for 44 hours. There was obtained 290 mg product. 54 % yield.

IR(CHCl₃) 1693 cm⁻¹. ¹H NMR: δ 0.86 (t, 3H); 1.23 (c, 12H); 1.41 (c, 2H); 1.78 (c, 2H); 3.02 (d, 2H); 3.5 (d, 2H); 6.7 (c, 3H); 7.18 (c, 4H).

EXAMPLE 68

N-(2,4,6-Trimethoxy)phenyl-2-[(2-(6-ethoxybenzthiazolyl)thio]octanamide

The title compound was prepared according to the procedures described in Examples 5 and 6.

M.p. 78-81 °C. ¹H NMR δ (CDCl₃) 8.13 (s, 1); 7.69 (d, 4 Hz, 1); 7.22 (d, 1 Hz, 1); 6.98 (dd, 1); 6.06 (s, 1); 4.65 (t, 4 Hz, 1); 4.07 (q, 3 Hz, 2); 3.81 (s, 2); 3.76 (s, 2); 3.54 (s, 6); 2.2-1.2 (m, 12); 0.89 (m, 3). IR (CHCl₃) 3400, 2929, 1690, 1520 cm⁻¹.

EXAMPLE 69

N-(2,4,6-Trimethoxy)phenyl-2-[(1-tert-nonyl)thio]octanamide

The title compound was prepared according to the procedures described in Examples 5 and 6.

Oil. ¹H NMR (CDCl₃) δ 8.10 (bs, 1); 3.82 (s, 3); 3.79 (s, 6); 3.40 (bs, 1); 1.98-0.80 (m, 32). IR(CHCl₃)

3320, 2950, 1670, 1600 cm^{-1} .

EXAMPLE 70

5

N-(5-Isoquinoliny)-2-((1-N-hexyl)thio)heptanamide

10 The title compound was prepared according to the procedures described in Examples 1 through 4.
M.p. 83-85 °C. ^1H NMR (CDCl_3) δ 9.30 (bs, 1); 9.16 (s, 1); 8.48 (d, 4 Hz, 1); 8.23 (d, 4 Hz, 1); 7.71-7.46 (m, 3); 3.47 (t, 2 Hz, 1); 2.58 (t, 5 Hz, 2); 2.08-0.76 (m, 20). IR (CHCl_3) 3317, 2921, 1682, 1515 cm^{-1} .

15

EXAMPLE 71

N-(2,4,6-Trimethoxy)phenyl-2-((1-N-hexyl)thio)decanoamide

20

The title compound was prepared according to the procedures described in Examples 5 and 6.
M.p. 55-56 °C. C, calc'd, 67.19; Found, 66.41. ^1H NMR (CDCl_3) δ 7.81 (bs, 1); 6.14 (s, 2); 3.81 (s, 3); 3.78 (s, 6); 3.48 (s, 6); 3.46 (t, Hz, 1); 2.69 (bs, 2); 1.98-0.76 (m, 28). IR (CHCl_3) 3340, 2930, 1670, 1600 cm^{-1} .

25

EXAMPLE 72

30

N-(2,4,6-Trimethoxy)phenyl-2-((1-n-hexyl)sulfonyl)octanamide

The title compound was prepared by treating N-(2,4,6-trimethoxy)phenyl-2-((1-n-hexyl)thio)octanamide (0.5 g, 1.18 mmol) with m-chloroperoxybenzoic acid (0.56 g, 2.76 mmol) in dichloromethane (10 mL).
35 Extractive work up and chromatography on silica gel (1:3 ethyl acetate:hexanes, eluent) provided N-(2,4,6-trimethoxy)phenyl-2-((1-n-hexyl)sulfonyl)octanamide (0.28 g, 52%).

M.p. 134-137 °C. ^1H NMR (CDCl_3) δ 7.44 (s, 1); 6.17 (s, 2); 3.85 (s, 3); 3.27 (m, 1); 2.20 (m, 1); 2.20-0.93 (m, 26). IR (CHCl_3) 3370, 2930, 1690, 1600 cm^{-1} .

40

EXAMPLE 73

N-(2,4,6-Trimethoxy)phenyl-2-((1-N-hexyl)thio)hexanamide

The title compound was prepared by a procedure similar to that described in Examples 1 through 4.
M.p. 67-69 °C. ^1H NMR (CDCl_3) δ 7.76 (bs, 1); 6.12 (s, 2); 3.78 (s, 3); 3.76 (s, 3); 3.42 (t, 4 Hz, 1); 2.69-2.60 (m, 2); 1.99-1.19 (m, 14); 0.92-0.79 (m, 6).

50

EXAMPLE 74

55

N-(2,4,6-Trimethoxy)phenyl-2-((1-N-pentyl)thio)octanamide

The title compound was prepared according to the procedures described in Examples 5 and 6.

M.p. 85-86 °C. ¹H NMR δ (CDCl₃) 7.78 (s, 1); 6.11 (s, 2); 3.77 (s, 3); 3.74 (s, 6); 3.44 (t, 4 Hz, 1); 2.66 (m, 2); 1.90-1.22 (m, 16); 0.87 (m, 6). IR (CHCl₃) 3350, 2930, 1675, 1610 cm⁻¹.

5

EXAMPLE 75

10

N-(2,4,6-Trimethoxy)phenyl-2-((1-N-hexyl)thio)pentanamide

The title compound was prepared according to the procedures described in Examples 5 and 6.

M.p. 74-75 °C. ¹H NMR (CDCl₃) δ 7.81 (bs, 1); 6.15 (s, 2); 3.82 (s, 3); 3.79 (s, 6); 3.48 (t, 3 Hz, 1); 2.70 (m, 2); 1.98-0.85 (m, 8). IR (CHCl₃) 3320, 2920, 1675, 1600 cm⁻¹.

15

EXAMPLE 76

20

N-(2,4,6-Trimethoxy)phenyl-2-((1-N-hexyl)thio)heptanamide

The title compound was prepared according to the procedures described in Examples 5 and 6.

M.p. 89-91 °C. ¹H NMR (CDCl₃) δ 7.82 (bs, 1); 6.16 (s, 2); 3.82 (s, 3); 3.80 (s, 6); 3.48 (t, 3 Hz, 1); 2.70 (m, 2); 2.00-0.84 (m, 22). IR (CHCl₃) 3340, 2930, 1675, 1600 cm⁻¹.

25

EXAMPLE 77 AND 78

30

N-(2,4,6-Trimethoxy)phenyl-2-((1-N-hexyl)sulfinyl)octanamide (Diastereomer A and Diastereomer B)

The title compound was prepared by treating N-(2,4,6-trimethoxy)phenyl-2-(1-n-hexyl)thiooctanamide (0.5 g, 1.2 mmol) with m-chloroperoxybenzoic acid (0.29 g, 1.4 mmol) in dichloromethane (10 mL) at -20°C. Extractive work up and chromatography on silica gel (1:1 ethyl acetate:hexanes, eluent) provided Diastereomer A, m.p. 116-118 °C, (less polar, 0.18 g, 33%) and Diastereomer B, m.p. 105-106 °C, (more polar, 0.12 g, 12%).

35

40

EXAMPLE 79

45

N-(2,4,6-Trimethoxy)phenyl-2-((1-N-hexyl)thio)butanamide

The title compound was prepared according to the procedures described in Examples 5 and 6.

M.p. 79-80 °C. ¹H NMR (CDCl₃) δ 7.84 (bs, 1); 6.17 (s, 2); 3.82 (s, 2); 3.81 (s, 6); 3.44 (t, 3 Hz, 1); 2.71 (m, 2); 2.02-0.86 (m, 16). IR (CHCl₃) 3340, 2960, 1675, 1600 cm⁻¹.

50

EXAMPLE 80

55

N-(2,4,6-Trimethoxy)phenyl-2-((1-N-butyl)thio)octanamide

The title compound was prepared according to the procedures described in Examples 5 and 6.

M.p. 87-88 °C. ¹H NMR (CDCl₃) δ 7.74 (s, 1); 6.80 (s, 2); 3.75 (s, 3); (s, 6); 3.40 (t, 4 Hz, 1); 2.63 (m, 2);

2.04-1.19 (m, 14); 0.88-0.80 (m, 6). IR (CHCl₃) 2930, 1680, 1605 cm⁻¹.

EXAMPLE 81

5

N-(2,4,6-Trimethoxy)phenyl-2-((2-thiazolyl)thio)octamide

10 The title compound was prepared according to the procedures described in Examples 5 and 6.

M.p. 81-83 °C. ¹H NMR (CDCl₃) δ 8.10 (s, 1); 7.62 (m, 1); 7.19 (m, 1); 6.06 (s, 2); 4.40 (t, 3 Hz, 1); 3.75 (s, 3), 3.66 (s, 6); 2.2-1.22 (m, 10); 0.86 (m, 3). IR (CHCl₃) 2920, 1685, 1600, 1460 cm⁻¹.

The title compounds of Examples 82-96 were prepared by a procedure similar to that of Examples 5 and 6.

15

EXAMPLE 82

20

N-(2,4,6-Trimethoxy)-2-((benzthiazol-2-yl)thio)octanamide

M.p. 108-112 °C. ¹H NMR (CDCl₃) δ 8.26 (s, 1); 7.78 (d, 4 Hz, 1); 7.72 (d, 4 Hz, 1); 7.35 (m, 1); 6.09 (s, 1); 6.00 (s, 1); 4.98 (t, 4 Hz, 1); 3.76 (s, 3); 3.70 (s, 3); 3.46 (s, 4); 2.2-1.2 (m, 10); 0.84 (m, 3). IR (CHCl₃) 2930, 1690, 1600, 1510, 1465, 1440 cm⁻¹. N Calc.: 590 Found: 5.29.

25

EXAMPLE 83

30

N-(2,4,6-Trimethoxy)phenyl-2-((1-N-decyl)thio)butanamide

M.p. 58-59 °C. ¹H NMR (CDCl₃) δ 7.85 (bs, 1); 6.17 (s, 2); 3.83 (s, 3); 3.80 (s, 3); 3.48 (t, 3 Hz, 1); 2.71 (m, 2); 2.06-0.86 (m, 24). IR (CHCl₃) 3350, 2910, 1875, 1600 cm⁻¹.

35

EXAMPLE 84

40

N-(N-Pentyl)-2-(2-(6-ethoxybenzthiazolyl)thio)decanamide

M.p. 48-50 °C. ¹H NMR (CDCl₃) δ 7.68 (bs, 5 Hz, 1); 7.24 (bs, 1); 7.18 (d, 1 Hz, 1); 6.98 (q, 1 Hz, 1); 4.31 (t, 3 Hz, 1); 4.04 (q, 3, 6 Hz, 2); 3.22 (m, 2), 2.16-0.68 (m, 29). IR (CHCl₃) 3310, 2930, 1675, 1610 cm⁻¹.

45

EXAMPLE 85

50

N-(2,4,6-Trifluoro)phenyl-2-(N-hexyl)thiooctanamide

M.p. 63-66 °C. ¹H NMR (CDCl₃) δ 8.08 (bs, 1); 6.70 (t, 4 Hz, 2); 3.42 (t, 3 Hz, 1); 2.58 (t, 4 Hz, 2); 1.98-0.80 (m, 24). IR (CHCl₃) 3666, 29.23, 1692, 1644, 1511, 14.76 cm⁻¹.

55

EXAMPLE 865 N-(2,4,6-Trimethoxy)phenyl-2-(2-(6-ethoxybenzthiazolyl)thio)octanamide

Amorphous. ^1H NMR (CDCl_3) δ 8.24 (s, 1); 7.64 (d, 6 Hz, 1); 7.16 (d, 1 Hz, 1); 6.92 (m, 1); 6.11 (s, 1); 6.08 (s, 1); 6.00 (s, 1); 4.58 (t, 4 Hz, 1); 4.00 (q, 3 Hz, 2); 3.78 (s, 3); 3.74 (s, 2) 3.70 (s, 2); 3.46 (s, 3); 2.2-1.2 (m, 9); 0.85 (m, 3 Hz, 3).

10

EXAMPLE 87

15

N-(2,4,6-Trimethoxy)phenyl-2-(2-(6-ethoxybenzthiazolyl)thio)pentanoamide

M.p. 44-46 °C. ^1H NMR (CDCl_3) δ 8.18 (s, 1); 7.56 (d, 5 Hz, 1); 7.07 (d, 1 Hz, 1); 6.83 (m, 1); 5.91 (s, 2); 4.52 (t, 3 Hz, 1); 3.91 (q, 4 Hz, 2); 3.60 (s, 3); 3.38 (s, 6); 2.10-1.10 (m, 7); 0.88 (m, 3 Hz, 3). IR (CHCl_3) 2960, 1690, 1605, 1510, 1470 cm^{-1} .

20

EXAMPLE 88

25

N-(2,4,6-Trimethoxy)phenyl-2-(2-(5-chlorobenzthiazolyl)thio)hexanamide

M.p. 100-101 °C. ^1H NMR (CDCl_3) δ 8.20 (s, 1); 7.65 (d, 4 Hz, 1); 7.20 (m, 1); 7.08 (s, 1); 6.10 (s, 2); 4.72 (t, 4 Hz, 1); 3.79 (s, 9); 2.2-1.1 (m, 10); 0.88 (m, 3 Hz, 3).

30

EXAMPLE 89

35

N-(2,4,6-Trimethoxy)phenyl-2-(2-(6-ethoxybenzthiazolyl)thio)hexanamide

Amorphous. ^1H NMR (CDCl_3) δ 8.28 (s, 1); 7.66 (d, 5 Hz, 1); 6.94 (m, 1); 6.02 (s, 2); 4.61 (t, 4 Hz, 1); 4.03 (q, 3, Hz, 2); 3.74 (s, 3); 3.50 (s, 6); 2.15 (s, 2); 1.96-1.32 (m, 7); 0.92 (t, 4 Hz, 3). IR (CHCl_3) 2930, 1690, 1600, 1510, 1450 cm^{-1} .

40

EXAMPLE 90

45

N-(5-Isoquinolinyl)-2-((1-tet-nonyl)thio)hexanamide

M.p. 83-85. ^1H NMR (CDCl_3) δ 9.4 (s, 1); 9.2 (s, 1); 8.4 (m, 1), 7.8 (m, 1); 7.6 (m, 2); 3.55 (m, 1); 2.65 (m, 2); 2.2-1.8 (m, 2); 1.0-1.80 (m, 20); 0.8-1.0 (m, 6). IR (CHCl_3) 2920, 2860, 1690, 1530 cm^{-1} .

50

EXAMPLE 91

55

N-(5-Isoquinolinyl)-2-((1-N-butyl)thio)octanamide

Amorphous. ^1H NMR (d-DMSO) δ 9.42 (s, 1); 9.30 (m, 1); 8.60 (m, 1), 8.40 (m, 1); 7.82 (m, 2); 7.65 (m, 2); 3.60 (m, 1); 2.65 (m, 2); 2.1 (m, 1); 1.90 (m, 1); 1.2-1.8 (m, 15); 0.9 (m, 6). IR (CHCl_3) 2950, 2860, 1690, 1520, 1480 cm^{-1} .

5

EXAMPLE 9210 N-(5-Isoquinolinyl)-2-(2-(6-ethoxybenzthiazolyl)thio)octanamide

M.p. 92-93° C. ^1H NMR (CDCl_3) δ 9.13 (s, 1); 8.36 (d, 5 Hz, 1); 8.30 (d, 3 Hz, 1), 7.78 (d, 6 Hz, 1); 7.70 (t, 5 Hz, 3); 7.53 (t, 4 Hz, 1); 7.19 (s, 1); 7.0 (m, 1); 4.61 (t, 4 Hz, 1); 4.03 (q, 3 Hz, 2); 2.3-1.8 (m, 2); 1.6-1.2 (m, 11); 0.82 (m, 3). IR (CHCl_3) 2920, 1700, 1610, 1550, 1470 cm^{-1} .

15

EXAMPLE 93

20

N-(2,4,6-Trifluorophenyl)-2-(2-(6-ethoxybenzthiazolyl)thio)decanamide

Amorphous. ^1H NMR (CDCl_3) δ 8.54 (s, 1); 7.67 (d, 4 Hz, 1); 7.20 (d, 1 Hz, 1); 7.00 (m, 1); 6.8-6.6 (m, 2); 4.48 (t, 3 Hz, 1); 4.03 (q, 3 Hz, 2); 2.2-1.0 (m, 17); 0.84 (m, 3). IR (CHCl_3) 2920, 2840, 1700, 1600, 1520 cm^{-1} .

25

EXAMPLE 94

30

N-(2,4,6-Trimethoxyphenyl)-2-(2-(6-ethoxybenzthiazolyl)thio)tetradodecanamide

M.p. 87-89° C. ^1H NMR (CDCl_3) δ 8.27 (s, 1); 7.68 (d, 5 Hz, 1); 7.20 (d, 1 Hz, 1); 6.96 (m, 1); 6.04 (s, 2); 4.62 (t, 3 Hz, 1); 4.05 (q, 3 Hz, 2); 3.74 (s, 3); 3.51 (s, 6); 2.2-1.2 (m, 25); 0.84 (m, 3). IR (CHCl_3) 2920, 2860, 1690, 1600, 1510 cm^{-1} .

35

EXAMPLE 95

40

N-(6-Methoxyisoquinolin-5-yl)-2-((1-n-hexyl)thio)decanamide

Amorph. ^1H NMR (CDCl_3) δ 9.1 (s, 1); 8.5 (m, 1); 8.4 (m, 1); 7.9 (m, 1); 7.5 (m, 2); 7.35 (m, 2); 4.0 (s, 3); 3.5 (m, 1); 2.7 (m, 2); 2.05 (m, 1); 1.85 (m, 1); 1.2-1.8 (m, 15); 0.9 (m, 6). IR (CHCl_3) 2920, 1680, 1625, 1485 cm^{-1} .

45

50

EXAMPLE 96N-(5-Isoquinolinyl)-2-((1-n-butyl)thio)decanamide

55

M.p. 73-75° C. ^1H NMR (CDCl_3) δ 9.8 (s, 1); 9.5 (m, 1); 8.6 (m, 1), 8.35 (m, 1); 7.8 (m, 1); 7.5 (m, 2); 3.5 (m, 1); 2.65 (m, 2); 2.1-1.7 (m, 18); 0.9 (m, 6). IR (CHCl_3) 2920, 1685, 1520, 1480 cm^{-1} .

EXAMPLE 975 N-(2,4,6-trimethoxyphenyl)-N'-[4-(3-methylbutyl)phenylmethyl]-N'-heptylurea

209 mg (1 mmole) 2,4,6-trimethoxyphenylisocyanate, 275 mg 4-(3-methylbutyl)benzylamine, and 10 ml methylene chloride were stirred at room temperature overnight. The reaction mixture was concentrated in vacuo. Chromatography on 100 g silica gel eluting with 1:1 hexane-ethyl acetate gave 320 mg product.

10 66% yield. ¹H NMR (CDCl₃): δ 0.81-0.96 (c) and 0.92 (d) (total 9H); 1.27 (c, 9H); 1.44-1.68 (c, 4H); 2.6 (t, 2H); 3.33 (t, 2H); 3.75 (s, 6H); 3.77 (s, 3H); 4.55 (s, 2H); 5.55 (s, 1H); 6.11 (s, 2H); 7.15 (d, 2H); 7.24 (d, 2H). IR (CHCl₃): 1654 cm⁻¹.

15 EXAMPLE 9820 N-(2,4,6-trimethoxyphenyl)-N'-[4-(2,2-dimethylpropyl)phenylmethyl]-N-heptylurea

The title compound was prepared according to the procedure of Example 97, but using 760 mg (3.63 mmole) 2,4,6-trimethoxyphenylisocyanate, 1.0 g (3.63 mmole) 4-(2,2-dimethylpropyl)benzylamine, and 20 ml methylene chloride. There was obtained 1.25 g product.

25 71% yield. ¹H NMR (CDCl₃): δ 0.82-0.95 (c) and 0.89 (s) (total 12H); 1.27 (c, 8H); 1.61 (c, 2H); 2.48 (s, 2H); 3.34 (t, 2H); 3.76 (s, 6H); 3.78 (s, 3H); 4.57 (s, 2H); 5.59 (s, 1H); 6.12 (s, 2H); 7.1 (d, 2H); 7.23 (d, 2H). IR (CHCl₃): 1657 cm⁻¹.

30 EXAMPLE 99

N-(6-Methylthio-8-acetaminoquinolin-5-yl)-2-(hexylthio) decanoic amide

35 5-Amino-6-methylthio-8-acetaminoquinoline, prepared according to the procedure of Gilman et al. (J. Amm. Chem. Soc. 68, 1577 (1946)), was coupled with 2-hexanethiodecanoic acid (prepared as described in Example 25) using the procedure described in Example 25, to give the title compound.

¹H NMR (CDCl₃): δ 9.75 (s, 1H); 8.82 (s, 1H); 8.68 (d, 5Hz, 1H); 8.46 (s, 1H); 7.97 (d, 7Hz, 1H); 7.41 (dd, 5 & 7Hz, 1H); 3.50 (t, 6Hz, 1H); 2.79 (t, 6Hz, 2H); 2.58 (s, 3H); 2.35 (s, 3H); 2.13 (m, 1H); 1.85 (m, 1H); 1.76-40 1.22 (m, 20H); 0.86 (m, 6H).

¹³C NMR (CDCl₃): 172.1, 169.0, 147.1, 136.4, 136.1, 134.5, 131.6, 125.2, 122.3, 113.1, 59.8, 51.1, 33.1, 32.3, 31.9, 31.4, 29.4, 29.3, 28.6, 27.8, 25.1, 22.6, 22.5, 15.3, 14.1.

IR (KBr): 3240, 2920, 1640, 1650, 1530 cm⁻¹.

45 The title compounds of Examples 100-107 were prepared according to the procedure described in Example 4.

EXAMPLE 10050 2-(4-t-Butylphenylthio)-N-(2,4,6-trimethoxyphenyl)octanamide

IR (CHCl₃): 1670 cm⁻¹.

55 EXAMPLE 101

2-(4-t-Butylphenylthio)-N-(2,4,6-trimethylphenyl)octanamide

IR (CHCl₃): 1670 cm⁻¹.

5

EXAMPLE 102

10 2-[4-(1,1-Dimethylpropyl)phenylthio]-N-(2,4,6-trimethylphenyl)nonanamide

IR (CHCl₃): 1670 cm⁻¹.

15

EXAMPLE 103

2-(4-n-Butylphenylthio)-N-(2,4,6-trimethylphenyl)nonanamide

20

IR (CHCl₃): 1669 cm⁻¹.

25

EXAMPLE 104

2-[4-(1,1-Dimethylpropyl)phenoxy]-N-(2,4,6-trimethoxyphenyl)octanamide

30

IR (CHCl₃): 1681 cm⁻¹.

35

EXAMPLE 105

2-[4-(1,1-Dimethylpropyl)phenoxy]-N-(2,4,6-trimethylphenyl)octanamide

40

IR (CHCl₃): 1678 cm⁻¹.

45

2-(4-n-Propylphenoxy)-N-(2,4,6-trimethylphenyl)decanamide

IR (CHCl₃): 1678 cm⁻¹.

50

EXAMPLE 107

55 2-(4-n-Propylphenylthio)-N-(2,4,6-trimethylphenyl)-nonanamide

IR (CHCl₃): 1669 cm⁻¹.

The title compounds of Examples 108-120 were prepared according to the procedure described in

Examples 5 and 6.

EXAMPLE 108

5

N-(2,4,6-Trimethyl)phenyl-2-((2-methylfuryl)thio)octanamide

10 M.p. 73-75 ° C.

EXAMPLE 109

15

N-(2,4,6-Trimethyl)phenyl-2-((2-benzimidazolyl)thio)decaneamide

M.p. 171-172 ° C.

20

EXAMPLE 110

25

N-(2,4,6-Trimethyl)phenyl-2-((2-benzothiazolyl)thio)octanamide

M.p. 103-106 ° C.

30

EXAMPLE 111

35 N-(2,4,6-Trimethyl)phenyl-2-(1-hexylthio) octanamide

M.p. 69-72 ° C.

40

EXAMPLE 112

N-(2,4,6-Trimethyl)phenyl-2-[2-(3-hydroxy-2-pyridyl)thio] decanoamide

45

IR (CHCl₃): 3200, 2920, 1675 cm⁻¹.

EXAMPLE 113

50

N-(2,4,6-Trimethyl)phenyl-2-[2-(6-chlorobenzothiazolyl)thio] octamide

55 M.p. 130-131 ° C.

EXAMPLE 114

N-(2,4,6-Trimethyl)phenyl-2-(1-heptylthio) octanamide

5

M.p. 53-56 ° C.

EXAMPLE 115

10

N-(Isoquinolin-5-yl)-2-(1-heptylthio) decanamide

15

M.p. 71 ° C.

EXAMPLE 116

20

N-(2,4,6-Trimethyl)phenyl-2-(tert-butylthio) octanamide

M.p. 145-147 ° C.

25

EXAMPLE 117

30

N-(Isoquinolin-5-yl)-2-(4-propylphenylthio)decanoic amide

M.p. 86-88 ° C.

35

EXAMPLE 118

N-(Isoquinolin-5-yl)-2-(phenylmethylthio)decanoic amide

M.P. 86-88 ° C.

45

EXAMPLE 119

N-(Isoquinolin-5-yl)-2-(cyclohexylthio)decanoic amide

50

M.p. 98-100 ° C.

EXAMPLE 120

55

N-(Quinolin-5-yl)-2-(hexylthio)decanoic amide

IR (KBr) cm^{-1} : 3240, 2920, 2850, 1657, 1529.

The title compounds of Examples 121-122 were prepared according to the procedure described in Example 31.

5

EXAMPLE 121

10 N-(6-Methylquinolin-5-yl)-2-(hexylthio)decanoic amide

Mass spectrum m/e (relative intensity): M^+ 428.26 (1), 312.22 (23), 213.06 (30), 200.10 (23), 158.06 (100). High resolution mass spectra: m/e 428.2843, calc'd for $\text{C}_{26}\text{H}_{40}\text{N}_2\text{OS}$: 428.2853. Anal.: Calc'd for $\text{C}_{26}\text{H}_{40}\text{N}_2\text{OS}$: C, 72.85; H, 9.41; N, 6.54. Found: C, 73.04; H, 9.20; N, 6.52.

15

EXAMPLE 122

20

N-(4-Methoxycarbonyl-6-methoxyquinolin-5-yl)-2-(hexylthio)decanoic amide

IR (CHCl_3) cm^{-1} : 3320, 2915, 2862, 1748, 1651.

25

EXAMPLE 123

30 N-(Quinolin-5-yl)-2-(decyl)cyclopentane carboxamide

5-Aminoquinoline was converted to the title compound according to the procedure described in Example 30.

M.p. 75°C .

35

EXAMPLE 124

40

N-(6-Methoxyquinolin-5-yl)-2-(decyl)cyclopentane carboxamide

6-Methoxy-5-aminoquinoline, prepared according to the procedure of Example 33, was converted to the title compound according to the procedure described in Example 30.

M.p. $57-58^\circ\text{C}$.

45

EXAMPLE 125

50

N-(6-Methoxyquinolin-5-yl)-2-(4-sec-butylphenoxy)nonanoic amide

The title compound was prepared according to the procedure described in Example 4.

Anal: calc'd for $\text{C}_{29}\text{H}_{38}\text{N}_2\text{O}_3$: C, 75.35; H, 8.28; N, 6.06. Found: C, 74.81; H, 8.24; N, 5.96.

55

EXAMPLE 126

N-(6-Methoxyquinolin-5-yl)-2-octanyl-1,3-dithiane-2-carboxamide

5 5-Amino-6-methoxyquinoline, prepared as described in Example 60, was coupled with 2-octanyl-1,3-dithiane-2-carboxylic acid, prepared by treatment of 1,3-dithiane-2-carboxylic acid with sodium hexamethyl-

10 Oil. ^1H NMR(CDCl_3): δ 9.00 (s, 1H); 8.63 (d, 4Hz, 1H); 7.94 (d, 10Hz, 1H); 7.92 (d, 9Hz, 1H); 7.73 (d, 10Hz, 1H); 7.13 (dd, 4 & 12Hz, 1H); 3.94 (s, 3H); 3.10 (dt, 2 & 12Hz, 2H); 2.70 (dt, 4 & 12Hz, 2H); 2.07 (m, 2H); 1.96 (m, 2H); 1.60 (m, 2H); 1.31-1.08 (m, 10H); 0.87 (t, 6Hz, 3H).

The title compounds of Examples 127-131 were prepared according to the procedure described in Example 35.

EXAMPLE 127N-(6-Cyclohexylthio)quinolin-5-yl)-2-hexylthiodecanoic amide

M.p. 88-89 °C.

EXAMPLE 128N-(6-(3-Phenylpropylthio)quinolin-5-yl)-2-hexylthiodecanoic amide

M.p. 63-64 °C.

EXAMPLE 129N-(6-(benzylthio)quinolin-5-yl)-2-hexylthiodecanoic amide

40 Anal: calc'd for $\text{C}_{32}\text{H}_{44}\text{N}_2\text{OS}_2$: C, 71.48; H, 8.50; N, 5.20. Found: C, 71.59; H, 8.26; N, 5.21.

EXAMPLE 130N-(6-hexylthio)quinolin-5-yl)-2-hexylthiodecanoic amide

50 Anal: calc'd for $\text{C}_{31}\text{H}_{50}\text{N}_2\text{OS}_2$: C, 70.13; H, 9.49; N, 5.28. Found: C, 69.99; H, 9.37; N, 5.42.

EXAMPLE 131N-(6-chloroquinolin-5-yl)-2-hexylthiodecanoic amide

Anal: calc'd for $C_{25}N_3ClN_2OS$: C, 66.06; H, 8.7; N, 5.42. Found: C, 66.86; H, 8.31; N, 6.24.

The title compounds of Examples 132-141 were prepared according to the procedure described in Examples 54-59.

5

EXAMPLE 132

10 Exo-3-n-nonyl-endo-N-(2,4,6-trimethylphenyl)-bicyclo[2.2.1]hept-5-ene-2-carboxamide

IR ($CHCl_3$): 1659 cm^{-1} .

15

EXAMPLE 133

20 Exo-3-n-heptyl-endo-N-(2,4,6-trimethylphenyl)-bicyclo[2.2.1]hept-5-ene-2-carboxamide

IR ($CHCl_3$): 1660 cm^{-1} .

25

EXAMPLE 134

30 Exo-3-n-octyl-endo-N-(2,4,6-trimethoxyphenyl)bicyclo[2.2.1]hept-5-ene-2-carboxamide

IR ($CHCl_3$): 1660 cm^{-1} .

35

EXAMPLE 135

40 Endo-3-n-octyl-exo-N-(2,4,6-trimethoxyphenyl)bicyclo[2.2.1]hept-5-ene-2-carboxamide

IR ($CHCl_3$): 1677 cm^{-1} .

45

EXAMPLE 136

50 Endo-3-n-heptyl-exo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.1]hept-5-ene-2-carboxamide

IR ($CHCl_3$): 1670 cm^{-1} .

55

EXAMPLE 137

60 Exo-3-n-heptyl-endo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.1] hept-5-ene-2-carboxamide

IR ($CHCl_3$) 1660 cm^{-1} .

EXAMPLE 138

5 Exo-3-n-heptyl-endo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.2]oct-5-ene-2-carboxamide

IR (CHCl₃): 1664 cm⁻¹.

10

EXAMPLE 139

Endo-3-n-heptyl-exo-N-(2,4,6-trimethoxyphenyl)bicyclo[2.2.2]oct-5-ene-2-carboxamide

15

IR (CHCl₃): 1680 cm⁻¹.

20

EXAMPLE 140

Exo-3-n-nonyl-endo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.2]oct-5-ene-2-carboxamide

25

IR (CHCl₃): 1666 cm⁻¹.

30

EXAMPLE 141

Endo-3-n-nonyl-exo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.2]oct-5-ene-2-carboxamide

35

IR (CHCl₃): 1679 cm⁻¹.

40

EXAMPLE 142

2-n-Nonyl-N-(2,4,6-trimethylphenyl)indane-2-carboxamide

The title compound was prepared according to the procedure described in Examples 60-64.

IR (CHCl₃): 1671 cm⁻¹.

45

The title compounds of Examples 143-144 were prepared according to the procedure described in Examples 55-59.

50

EXAMPLE 143

Exo-2-n-decyl-endo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.1]hept-5-ene-2-carboxamide

55

IR (CHCl₃) 1676 cm⁻¹.

EXAMPLE 144

Endo-2-n-decyl-exo-N-(2,4,6-trimethoxyphenyl)-bicyclo[2.2.1]hept-5-ene-2-carboxamide

5

IR 1675 cm⁻¹.

The title compounds of Examples 145-147 were prepared according to the procedure described in Examples 60-64.

10

EXAMPLE 14515 2-n-Octyl-N-(2,4,6-trimethylphenyl)-1,2,3,4-tetrahydro-2-naphthamide

¹H NMR: δ 0.87 (t, 3H); 1.29 (c, 10H) 1.52 (c, 3H); 1.89 (c) and 1.191 (s) (total 8H); 2.2 (s, 3H); 2.3 (c, 1H); 2.91 (c) and 2.94 (d) (total 3H); 3.19 (d, 2H); 6.76 (s, 1H); 6.79 (s, 2H); 7.12 (s, 4H).

20

EXAMPLE 14625 2-n-Decyl-N-(2,4,6-trimethylphenyl)indane-2-carboxamideIR (CHCl₃) 1671 cm⁻¹.

30

EXAMPLE 14735 2-n-Nonyl-N-(2,4,6-trimethylphenyl)-1,2,3,4-tetrahydro-2-naphthamideIR (CHCl₃) 1668 cm⁻¹.

The title compounds of Examples 148-150 were prepared according to the procedure described in Examples 5 and 6.

40

EXAMPLE 14845 N-(2,6-Difluoro)phenyl-2-((1-N-hexyl)thio)octanamide

M.p. 46-48° C.

50

EXAMPLE 149N-(2,6-Difluoro)phenyl 2-(2-(6-ethoxybenzthiazolyl)thio)octanamide

55

M.p. 110-112° C.

EXAMPLE 150

5 N-(2,6-Difluoro)phenyl-2-(2-(6-ethoxybenzthiazolyl)thio)decanamide

M.p. 99-100° C.

10

EXAMPLE 151

15 N-(6-acetaminoquinolin-5-yl)-2-(hexylthio)decanoic amide

15

Commercially available 5-amino-6-nitroquinoline was reduced to 5,6-diaminoquinoline using a procedure analogous to that described in Example 34, except tin (II) chloride was used in place of iron. 5,6-Diaminoquinoline was converted to 5-amino-6-acetaminoquinoline by reaction with acetic anhydride and pyridine. Using the procedure outlined in Example 25, 5-amino-6-acetaminoquinoline and 2-(hexylthio)-
20 decanoic acid were coupled to give the title compound. Mass spectrum m/e: 471.3 (M+).

¹H NMR, δ (CDCl₃): 8.99 (s, 1H); 8.85 (dd, J=2 & 4 Hz, 1H); 8.37 (s, 1H); 8.14 (d, J=10 Hz, 1H), 7.89 (d, J=9 Hz, 1H); 7.79 (d, J=9 Hz, 1H); 7.36 (dd, J=4 & 10 Hz, 1H); 3.47 (t, J=8 Hz, 1H); 2.64 (t, J=8 Hz, 2 H); 2.13 (s, 3H); 1.87 (m, 2H); 1.28 (br m, 20 H); 0.87 (m, 6H).

25

EXAMPLE 152

30 N-(6-aminoquinolin-5-yl)-2-(hexylthio)decanoic amide

N-(6-Acetaminoquinolin-5-yl)-2-(hexylthio)decanoic amide, prepared as described in Example 151, was treated with aqueous hydrochloric acid and isopropanol to give the title compound. Mass spectrum m/e: 428.3 (M+ - 1).

35

¹H NMR, δ (CDCl₃): 8.94 (d, J=6 Hz); 8.93 (d, J=2 Hz, 1H), 7.93 (d, J=9 Hz, 1H); 7.73 (d, J=9 Hz, 1H); 7.50 (dd, J=2 & 6 Hz, 1H); 4.33 (t, J=7 Hz, 1H); 2.43 (m, 2H), 2.00 (m, 2H); 1.48 (m, 4 H); 1.19 (m, 16H); 0.81 (m, 6 H).

40

EXAMPLE 153

45 N-(6-methylthioquinolin-5-yl)oleamide

45

5-Amino-6-methylthioquinoline, prepared as described in Example 34, and commercially available oleoyl chloride were coupled to give the title compound according to the procedure described in Example 25. Mass spectrum m/e: 454.3 (M+).

50

¹H NMR, δ (CDCl₃): 8.77 (d, J=3 Hz, 1 H) 8.00 (d, J=6 Hz, 1H); 7.98 (d, J=7 Hz, 1H) 7.59 (d, J=6 Hz, 1H); 7.32 (dd, J=3 & 7 Hz, 1H); 7.14 (s, 1H); 5.32 (br s, 2H); 2.51 (s, 5H) 1.99 (br s, 4H); 1.81 (m, 2H); 1.30 (br s, 20H); 0.84 (t, J=6 Hz, 3H).

55

EXAMPLE 154

N-(8-amino-6-methoxyquinolin-5-yl)-2-hexylthiodecanoic amide

Commercially available 6-methoxy-8-aminoquinoline (Chemical Procurement Laboratories) was acetylated with acetic anhydride and pyridine. The resultant 6-methoxy-8-acetaminoquinoline was nitrated and reduced using the procedure described in Example 33 to give 5-amino-6-methoxy-8-acetaminoquinoline. This product was coupled with 2-(hexylthio)decanoic acid according to Example 25 and hydrolysed with aqueous hydrochloric acid and isopropanol to give the title compound. Mass spectrum m/e: 459.3 (M+).

¹H NMR, δ (CDCl₃): 8.52 (d, J=4 Hz, 1H), 8.19 (s, 1H); 7.85 (d, J=8 Hz, 1H); 7.27 (dd, J=4 & 8 Hz, 1H); 6.60 (s, 1H); 5.06 (br s, 2H); 3.82 (s, 3H); 3.48 (t, J=6 Hz, 1H); 2.27 (t, J=7 Hz, 2H), 2.01 (m, 1H); 1.82 (m, 1H); 1.63 (m, 2H); 1.28 (m, 18H); 0.86 (m, 6H).

EXAMPLE 155

N-(6-(1,2,4-triazol-3-yl)thioquinolin-5-yl)-2-hexylthiodecanoic amide

Using a procedure analogous to that described in Example 34, 5-amino-6-(1,2,4-triazol-3-yl)quinoline was synthesized and coupled with 2-(hexylthio)decanoic acid to give the title compound.

Analysis: C 62.94, H 7.56, N 13.34; calc. for C₂₇H₃₃N₅OS₂: C 63.12, H 7.65, N 13.63.

¹H NMR, δ (CDCl₃): 9.59 (s, 1H), 8.89 (d, J=5 Hz, 1H); 8.18 (d, J=9 Hz, 1H); 8.06 (s, 1H); 7.93 (d, J=9 Hz, 1H); 7.65 (d, J=9 Hz, 1H), 7.43 (dd, J=5 & 9 Hz, 1H), 3.50 (t, J=7 Hz, 1H); 2.74 (t, J=7 Hz, 2H), 2.04 (m, 1H); 1.70 (m, 1H); 1.59 (m, 3H); 1.25 (m, 18H); 0.86 (m, 6H).

EXAMPLE 156

N-(6-methylthioquinolin-5-yl)-2,2-di(hexylthio)acetamide

2,2-Di(hexylthio)acetic acid was synthesized from dichloroacetic acid and hexanethiol using a procedure similar to that described in Example 1. 2,2-Di(hexylthio)acetic acid and 5-amino-6-methylthioquinoline were coupled to give the title compound using the procedure described in Example 25.

Mass spectrum m/e: 464.2 (M+).

¹H NMR δ (CDCl₃): 8.83 (dd, J=3 & 4 Hz, 1H); 8.45 (s, 1H), 8.12 (d, J=8 Hz, 1H); 8.09 (d, J=8 Hz, 1H); 7.64 (d, J=8 Hz, 1H); 7.42 (dd, J=4 & 8, 1H); 4.51 (s, 1H); 2.89 (t, J=8 Hz, 4H); 2.57 (s, 3H), 1.68 (m, 6H), 1.44 (m, 6H); 1.32 (m, 8H); 0.81 (t, J=7 Hz, 6H).

EXAMPLE 157

N-(6-methylthioquinolin-5-yl)-2-heptylnonanoic amide

By use of the procedures described in Example 25, nonanoic acid was alkylated with heptyl bromide and the resulting product was coupled with 5-amino-6-methylthioquinoline (Example 34) to give the title compound.

Mass spectrum m/e: 428.3 (M+ - SCH₃).

¹H NMR (CDCl₃): δ 8.77 (d, J=4 Hz, 1H); 8.00 (d, J=8 Hz, 1H); 7.95 (d, J=8 Hz, 1H); 7.55 (d, J=8 Hz, 1H); 7.49 (s, 1H); 7.31 (dd, J=4 & 8 Hz, 1H); 2.49 (s, 3H); 2.43 (m, 1H); 1.40 (br, 24H); 0.85 (t, J=6 Hz, 6H).

EXAMPLE 158

N-(6-methylthioquinolin-5-yl)-2-[2-(6-ethoxybenzthiazolyl)-thio]decanoic amide.

The title compound was prepared by procedures analogous to those as described in Examples 5 and 6.

MP: 139-141 °C

5 ¹H NMR (CDCl₃): δ 9.42 (s, 1H); 8.76 (d, 3Hz, 1H); 8.00 (d, 9Hz, 2H); 7.74 (d, 9Hz, 1H); 7.60 (d, 9Hz, 1H); 7.21 (m, 2H); 6.98 (dd, 3 & 9 Hz, 1H); 4.71 (t, 7Hz, 1H); 4.05 (q, 7 Hz, 2H); 2.53 (s, 3H); 2.30 (m, 1H); 1.97 (m, 1H); 1.59 (m, 2H); 1.45-1.25 (m, 10H), 1.00-0.81 (m, 6H).

FABMS m/e: 554 (M⁺ + 1)

Anal.: Calc'd for C₂₉H₃₅N₃O₂S₃ · 1/2 H₂O: C, 61.89; H, 6.45; N, 7.46.

10 Found: C, 62.14; H, 6.33; N, 7.43.

EXAMPLE 159

15

N-(3-methyl-6-chloro-8-acetaminoquinolin-5-yl)-2-(hexylthio) decanoic amide

3-Methyl-5-amino-6-chloro-8-acetaminoquinoline, prepared according to the procedures of Utermohlen, W.P., J. Org. Chem., 8, 544 (1943) and Gilman et al., J. Am. Chem. Soc., 68, 1577 (1946), was coupled with 2-hexanethiodecanoic acid (prepared as described in Example 25) using the procedure described in Example 25, to give the title compound.

MP: 140-141 °C

20 ¹H NMR (CDCl₃): δ 9.68 (s, 1H); 8.72 (s, 1H); 8.57 (s, 2H); 7.76 (s, 1H); 3.52 (t, 7Hz, 1H); 2.75 (t, 7Hz, 2H); 2.49 (s, 3H); 2.33 (s, 3H); 2.13-1.28 (m, 22H); 0.87 (m, 6H).

EIMS m/e: 519 (M⁺)

Anal.: Calc'd for C₂₈H₄₂N₃O₂SCl: C, 64.66; H, 8.14; N, 8.07.

Found: C, 64.65; H, 8.39; N, 7.96.

30

EXAMPLE 16035 N-(3-methyl-6-methylthio-8-acetaminoquinolin-5-yl)-2-(hexylthio) decanoic amide.

3-Methyl-5-amino-6-methylthio-8-acetaminoquinoline, prepared according to the procedures of Utermohlen, W.P., J. Org. Chem., 8, 544 (1943) and Gilman et al., J. Am. Chem. Soc., 68, 1577 (1946), was coupled with 2-hexanethiodecanoic acid (prepared as described in Example 25) using the procedure described in Example 25, to give the title compound.

40 MP: 128-131 °C

¹H NMR (CDCl₃): δ 9.75 (s, 1H); 8.76 (s, 1H); 8.54 (s, 1H); 8.38 (s, 1H); 7.74 (s, 1H); 3.52 (t, 7Hz, 1H); 2.80 (t, 7Hz, 2H); 2.58 (s, 3H); 2.48 (s, 3H); 2.34 (s, 3H); 2.15-1.22 (m, 22H); 0.87 (m, 6H).

EIMS m/e: 531 (M⁺)

45 Anal.: Calc'd for C₂₉H₄₅N₃O₂S₂: C, 65.50; H, 8.53; N, 7.90.

Found: C, 65.33; H, 8.55; N, 7.85.

EXAMPLE 161

50

N-(3-methyl-6-methylthioquinolin-5-yl)-2-(hexylthio)decanoic amide

55 3-Methyl-5-amino-6-methylthioquinoline, prepared according to the procedures of Utermohlen, W.P., J. Org. Chem., 8, 544 (1943) and Gilman et al., J. Am. Chem. Soc., 68, 1577 (1946), was coupled with 2-hexanethiodecanoic acid (prepared as described in Example 25) using the procedure described in Example 25, to give the title compound.

M.p.: 137-138 °C.

¹H NMR (CDCl₃): δ 8.69 (d, 2Hz, 1H) ; 8.52 (s, 1H); 8.02 (d, 9Hz, 1H) ; 7.79 (s, 1H); 7.58 (d, 9Hz, 1H); 3.54 (t, 7Hz, 1H); 2.81 (t, 7Hz, 2H); 2.55 (s, 3H); 2.49 (s, 3H); 2.15-1.25 (m, 22H); 0.87 (m, 6H).

EIMS m/e: 474 (M⁺)

5 Anal.: Calc'd for C₂₇H₄₂N₂O₅S₂; C, 68.31; H, 8.91; N, 5.90.

Found: C, 68.52; H, 8.94; N, 5.91.

EXAMPLE 162

10

N-(6-nitroquinolin-5-yl)-2-(hexylthio)decanoic amide.

15 Commercially available 5-amino-6-nitroquinoline was coupled with 2-hexanethiodecanoic acid (prepared as described in Example 25) using the procedure described in Example 25, to give the title compound.

M.p.: 89.91 °C.

¹H NMR (CDCl₃): δ 10.10 (s, 1H); 9.05 (dd, 2 & 4 Hz, 1H); 8.26 (d, 9 Hz, 1H); 8.24 (m, 1H); 8.09 (d, 9 Hz, 1H); 7.52 (dd, 4 & 9 Hz, 1H); 3.48 (dd, 6 & 8 Hz, 1H); 2.65 (m, 2H); 2.05 (m, 1H); 1.86 (m, 1H); 1.70-1.20 (m, 20H); 0.87 (t, 6 Hz, 6H).

20 FABMS m/e: 460 (M⁺ + H)

Anal.: Calc'd for C₂₅H₃₇N₃O₃S: C, 65.33; H, 8.11; N, 9.14.

Found: C, 65.42; H, 8.13; N, 9.23.

25

EXAMPLE 163

N-(6-N,N-dimethylaminoquinolin-5-yl)-2-(hexylthiol)decanoic amide

5-Nitro-6-chloroquinoline, prepared as described by Manske & Kulka, Organic Reactions, Vol. VII, 59 (1953) and Campbell et al., J. Am. Chem. Soc., 68, 1559 (1946), was allowed to react with dimethylamine to yield 5-nitro-6-dimethylaminoquinoline. This material was converted to the title compound by the procedure described in Example 33.

35 M.p.: oil

¹H NMR (CDCl₃): δ 8.80 (dd, 2 & 4 Hz, 1H); 8.69 (s, 1H); 8.02 (m, 1H); 8.00 (d, 9 Hz, 1H); 7.60 (d, 9 Hz, 1H); 7.35 (dd, 4 & 9 Hz, 1H); 3.50 (t, 8 Hz, 1H); 2.77 (s, 6H); 2.73 (t, 7 Hz, 2H); 2.12 (m, 1H); 1.86 (m, 1H); 1.72-1.25 (m, 20H); 0.87 (m, 6H).

40 FABMS m/e: 458 (M⁺ + H)

Anal.: Calc'd for C₂₇H₄₃NSO: C, 70.85; H, 9.47; N, 9.18;

Found: C, 70.59; H, 9.31; N, 9.10.

45

EXAMPLE 164

N-(6-trifluoromethylquinolin-5-yl)-2-(hexylthio)decanoic amide

50

5-Nitro-6-trifluoromethylquinoline, prepared as described by Manske & Kulka, Organic Reactions, Vol. VII, 59 (1953) and Campbell et al., J. Am. Chem. Soc., 68, 1559 (1946), was converted to the title compound by the procedure described in Example 33.

M.p.: oil

55 ¹H NMR (CDCl₃): δ 10.98 (s, 1H); 9.01 (d, 2Hz, 1H); 8.96 (dd, 2 & 4 Hz, 1H); 8.26 (dd, 2 & 8 Hz, 1H); 7.83 (s, 1H); 7.57 (dd, 4 & 8 Hz, 1H); 3.52 (t, 7 Hz, 1H); 2.61 (m, 2H); 2.02 (m, 1H); 1.84 (m, 1H); 1.65-1.15 (m, 20 H); 0.84 (t, 7 Hz, 3H); 0.78 (t, 7Hz, 3H).

FABMS m/e: 483 (M⁺ + H)

Anal.: Calc'd for $C_{26}H_{37}N_2SOF_3$: C, 64.70; H, 7.73; N, 5.80;
 Found: C, 64.37; H, 7.81; N, 5.71

5

EXAMPLE 165N-(cinnolin-5-yl)-2-(hexylthio)decanoic amide

10

The title compound was prepared by a procedure analogous to that described in Example 33.

M.p.: 54-56° C

1H NMR ($CDCl_3$): δ 9.41 (s, 1H); 9.36 (d, 6 Hz, 1H); 8.40 (d, 9Hz, 1H) ; 8.33 (d, 7 Hz, 1H); 7.86 (m, 2H); 3.54 (t, 6Hz, 1H); 2.63 (t, 7Hz, 2H); 2.03 (m, 1H); 1.81 (m, 1H); 1.68-1.20 (m, 20H); 0.84 (m, 6H).

15

HRFABMS m/e: 416.2805 ($C_{24}H_{37}N_2SO + H^+$ requires 416.2738)

EXAMPLE 166

20

N-(cinnolin-8-yl)-2-(hexylthiol)decanoic amide

The title compound was prepared by a procedure analogous to that described in Example 33.

25

M.p.: oil

1H NMR ($CDCl_3$): δ 11.10 (s, 1 H); 9.34 (d, 6Hz, 1H); 8.88 (dd, 1 & 8 Hz, 1H); 7.86 (d, 6Hz, 1H); 7.76 (t, 8Hz, 1H); 7.50 (dd, 1 & 8 Hz, 1H); 3.53 (t, 7Hz, 1H); 2.62 (t, 7Hz, 2H); 2.03 (m, 1H); 1.86 (m, 1H); 1.65-1.15 (m, 20H); 0.81 (m, 6H).

HREIMS m/e: 415.2648 ($C_{26}H_{37}N_2SO$ requires 415.2660)

30

EXAMPLE 167

35

N-(6-methylthiophthalazin-5-yl) oleic amide

5-Nitro-6-methylthiophthalazine, prepared as described by Sturrock et al. , Can. J. Chem. , 49 , 3047 (1971) and Hirsch & Orphanos, J. Heterocyclic Chem. , 2 , 206 (1965), was reduced with tin(II) chloride and HCl to yield 5-amino-6-methylthiophthalazine. This material was coupled with commercially available oleoyl chloride to yield the title compound.

40

M.p.: oil

1H NMR ($CDCl_3$): δ 9.34 (s,2H); 7.74 (d, 8Hz, 1H); 7.73 (s, 1H); 7.66 (d, 8Hz, 1H); 5.34 (t, 5Hz, 2H); 2.57 (t, 8 Hz, 2H); 2.54 (s, 3H); 2.01 (m, 4H); 1.81 (m, 2H); 1.50-1.10 (m, 20H); 0.86 (t, 6Hz, 3H).

45

FABMS m/e: 456 ($M^+ + 1$)

EXAMPLE 168

50

N-[2,4-bis(methylthio)pyridin-3-yl]-2-hexylthiodecanoic amide

The title compound was prepared in 13.2% yield according to the procedure of Example 4A.

55

1H NMR ($CDCl_3$): δ 0.86 (c, 6H); 1.17-1.76(c, 21H), 2.03 (m, 1H), 2.42 (s, 3H); 2.51(s, 3H); 2.77 (t, 2H); 3.46 (t, 1H); 6.82 (d, 1H), 8.23 (s, 1H); 8.26 (d, 1H).

IR ($CHCl_3$): 2920, 2851, 1679, 1553, 1465 cm^{-1} .

EXAMPLE 1695 N-[4,6-bis(methylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide

The title compound was prepared in 7% yield according to the procedure of Example 4.

¹H NMR (CDCl₃): δ 0.87 (c, 6H); 1.2-1.85 (c, 21 H); 2.02 (m, 1H); 2.52 (s, 6H); 2.74 (t, 2H); 3.45 (t, 1H); 8.18 (s, 1H); 8.65 (s, 1H).

10 IR (CHCl₃): 2923, 2852, 1681, 1521, 1466, 1406, 1357 cm⁻¹.

EXAMPLE 170

15

N-(6-methoxyisoquinolin-5-yl)-2-hexylthiodecanoic amide

The title compound was prepared in 62% yield according to the procedure of Example 4.

20 ¹H NMR (CDCl₃): δ 0.89 (c, 6H); 1.20-1.96 (c, 21 H); 2.07 (m, 1H); 2.75 (t, 2H); 3.55 (t, 1H); 4.0 (s, 3H); 7.39 (d, 1H); 7.48 (d, 1H); 7.94 (d, 1H); 8.45 (d, 1H); 8.52 (s, 1H); 9.14 (s, 1H).

IR (CHCl₃): 2922, 2852, 1674, 1624, 1465, 1380, 1323, 1267 cm⁻¹.

25

EXAMPLE 171N-(6-methoxyquinazolin-5-yl)-2-hexylthiodecanoic amide

30

The title compound was prepared in 15% yield according to the procedure of Example 4.

¹H NMR (CDCl₃): δ 0.88 (c, 6H); 1.18-1.94 (c, 21H); 2.08 (m, 1H); 2.71 (t, 2H); 3.55 (t, 1H); 4.0 (s, 3H); 7.69 (d, 1H); 8.0 (d, 1H); 8.78 (s, 1H); 9.21 (s, 1H); 9.31 (s, 1H).

IR (CHCl₃): 2923, 2852, 1682, 1621, 1573, 1496, 1476, 1465, 1372, 1319, 1273, 1255, 1222 cm⁻¹.

35

EXAMPLE 172

40

N-(4,6-dimethoxypyrimidin-5-yl)-2-hexylthiodecanoic amide

The title compound was prepared in 40% yield according to the procedure of Example 4.

45 ¹H NMR (CDCl₃): δ 0.87 (c, 6H); 1.22-2.0 (c, 22H); 2.64 (m, 1H); 3.43 (t, 1H); 3.97 (s, 6H); 7.90 (s, 1H); 8.33 (s, 1H).

IR (CHCl₃): 2922, 2852, 1680, 1582, 1491, 1465, 1410, 1399, 1312 cm⁻¹.

50

EXAMPLE 173N-(4,6-diethoxypyrimidin-5-yl)-2-hexylthiodecanoic amide

55

The title compound was prepared in 76% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (c, 6H); 1.19-1.70 (c, 27H); 1.82 (m, 1H); 2.64 (m, 2H); 3.45 (t, 1H); 4.39 (q, 4H); 7.89 (s, 1H); 8.28 (s, 1H).

IR (CHCl₃): 2924, 2853, 1681, 1582, 1491, 1441, 1386, 1315 cm⁻¹.

EXAMPLE 1745 N-[4-methoxy-6-(4-methoxyphenylthio)pyrimidin-5-yl]-2-hexyl- thiodecanoic amide

The title compound was prepared in 6% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (m, 6H); 1.17-2.04 (c, 22H); 2.72 (t, 2H); 3.50 (t, 1H); 3.83 (s, 3H); 3.96 (s, 3H); 6.94 (d, 2H); 7.44 (d, 2H); 8.17 (s, 1H); 8.37 (s, 1H).

10 IR (CHCl₃): 2900, 2840, 1700, 1600, 1565, 1480.

EXAMPLE 175

15

N-[4,6-bis(ethylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide

The title compound was prepared in 8% yield according to the procedure of Example 4B.

20 ¹H NMR (CDCl₃): δ 0.87 (m, 6H); 1.17-2.06 (c, 28H); 2.62 (m, 4H); 2.75 (t, 2H); 3.45 (t, 1H); 8.15 (s, 1H); 8.61 (s, 1H).

IR (CHCl₃): 2922, 2852, 1706, 1520, 1466, 1405, 1355 cm⁻¹.

25

EXAMPLE 176N-[4-methoxy-6-(2-ethoxyethylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide

30

The title compound was prepared in 38% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (m, 6H); 1.16-1.85 (c) and 1.19 (t) (total 24H); 1.94 (m, 1H); 2.68 (t, 2H); 3.32-3.57 (c), 3.52 (q) (total 5H); 3.65 (t, 2H); 3.95 (s, 3H); 8.03 (s, 1H); 8.47 (s, 1H).

35 IR (CHCl₃): 2952, 2925, 2854, 1684, 1562, 1541, 1481, 1408, 1385 cm⁻¹.

EXAMPLE 177

40

N-[2-(4-pyridinylthio)-4-methylpyridin-3-yl]-2-hexylthiodecanoic amide

The title compound was prepared in 10% yield according to the procedure of Example 4.

45 ¹H NMR (CDCl₃): δ 0.86 (m, 6H); 1.17-1.84 (c, 21H); 1.95 (m, 1H); 2.30 (s, 3H); 2.62 (t, 2H); 3.4 (t, 1H); 7.17 (d, 1H); 7.27 (m, 2H); 8.31 (d, 1H); 8.48 (b, 2H); 8.55 (s, 1H).

IR (CHCl₃): 2921, 2851, 1680, 1574, 1471 cm⁻¹.

EXAMPLE 178

50

N-[4-methoxy-6-(1-methyl-5-tetrazolythio)pyrimidin-5-yl]-2-hexylthiodecanoic amide

55 The title compound was prepared in 43% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (m, 6H); 1.18-1.87 (c, 21H); 1.98 (m, 1H); 2.65 (t, 2H); 3.49 (t, 1H); 4.02 (s, 3H); 4.12 (s, 3H); 8.26 (s, 1H); 8.58 (s, 1H).

IR (CHCl₃): 2900, 2840, 1690, 1560, 1485 cm⁻¹.

EXAMPLE 1795 N-[2-(2-furylmethylthio)-4-methylpyridin-3-yl]-2-hexylthiodecanoic amide

The title compound was prepared in 10% yield according to the procedure of Example 4B.

¹H NMR (CDCl₃): δ 0.87 (m, 6H); 1.17-2.03 (c, 22H); 2.19 (s, 3H); 2.65 (m, 2H); 3.42 (t, 1H); 4.47 (s, 2H); 6.24 (m, 2H); 6.92 (d, 1H); 7.30 (d, 1H); 8.18 (s, 1H); 8.25 (d, 1H).

10 IR (CHCl₃): 2920, 2850, 1706, 1675, 1481 cm⁻¹.

EXAMPLE 180

15

N-[2,4,6-tris(methylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide

The title compound was prepared in 79% yield according to the procedure of Example 4A.

20 ¹H NMR (CDCl₃): δ 0.87 (c, 6H); 1.17-1.86 (c, 21H); 2.01 (m, 1H); 2.50 (s, 6H); 2.56 (s, 3H); 2.73 (t, 2H); 3.43 (t, 1H); 8.06 (s, 1H).

IR (CHCl₃): 2922, 2852, 1686, 1499, 1465, 1347 cm⁻¹.

25

EXAMPLE 181N-(2,4,6-trimethoxypyrimidin-5-yl)-2-hexylthiodecanoic amide

30

The title compound was prepared in 74% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (m, 6H); 1.18-2.0 (c, 22H); 2.63 (m, 2H); 3.42 (t, 1H); 3.93 (s, 6H); 3.95 (s, 3H); 7.71 (s, 1H).

IR (CHCl₃): 2923, 2851, 1675, 1607, 1582, 1482, 1467, 1398, 1379 cm⁻¹.

35

EXAMPLE 182

40

N-[2-methyl-4,6-bis(ethylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide

The title compound was prepared in 52% yield according to the procedure of Example 4A.

45 ¹H NMR (CDCl₃): δ 0.87 (m, 6H); 1.19-1.84 (c, 27H); 2.0 (m, 1H); 2.57 (s, 3H); 2.75 (t, 2H); 3.15 (q, 4H); 3.44 (t, 1H); 8.04 (s, 1H).

IR (CHCl₃): 2920, 2852, 1680, 1467, 1406, 1359, 1314 cm⁻¹.

EXAMPLE 183

50

N-(6-methoxyquinolin-5-yl)-2-heptynonanoic amide

55

The title compound was prepared in 20% yield according to the procedure of Example 4.

¹H NMR (CDCl₃): δ 0.88 (m, 6H); 1.18-1.84 (c, 24H); 2.41 (m, 1H); 3.97 (s, 3H); 7.13 (s, 1H); 7.36 (q, 1H); 7.5 (d, 1H); 8.04 (t, 2H); 8.78 (m, 1H).

IR (CHCl₃): 2921, 2850, 1686, 1596, 1570, 1465, 1322, 1266 cm⁻¹.

EXAMPLE 1845 N-(2,4,6-trimethoxyphenyl)-2-heptylnonanoic amide

The title compound was prepared in 72% yield according to the procedure of Example 4.

¹H NMR (CDCl₃): δ 0.88 (m, 6H); 1.18-1.8 (c, 24H); 2.2 (m, 1H); 3.77 (s, 6H); 3.79 (s, 3H); 6.13 (s, 2H); 6.38 (s, 1H).

10 IR (CHCl₃): 2921, 2850, 1677, 1598, 1505, 1465, 1437, 1413, 1153, 1131 cm⁻¹.

EXAMPLE 185

15

N-(6-methoxyisoquinolin-5-yl)-2-heptylnonanoic amide

The title compound was prepared in 21% yield according to the procedure of Example 4.

20 ¹H NMR (CDCl₃): δ 0.88 (m, 6H); 1.18-1.85 (c, 24H); 2.41 (m, 1H); 3.98 (s, 3H); 7.09 (s, 1H); 7.37 (d, 1H); 7.52 (b, 1H); 7.91 (d, 1H); 8.44 (b, 1H); 9.13 (b, 1H).

IR (CHCl₃): 2922, 2850, 1685, 1625, 1465, 1381, 1324, 1279, 1268 cm⁻¹.

EXAMPLE 186

25

N-(4,6-dimethoxypyrimidin-5-yl)-2-heptylnonanoic amide

30

The title compound was prepared in 53% yield according to the procedure of Example 4.

¹H NMR (CDCl₃): δ 0.87 (m, 6H); 1.18-1.8 (c, 24H); 2.24 (m, 1H); 3.97 (s, 6H); 8.32 (s, 1H).

IR (CHCl₃): 2921, 2851, 1686, 1583, 1487, 1463, 1408, 1400, 1312, 1121 cm⁻¹.

35

EXAMPLE 18740 N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-heptylnonanoic amide

The title compound was prepared in 48% yield according to the procedure of Example 4B.

¹H NMR (CDCl₃): δ 0.87 (m, 6H); 1.17-1.82 (c, 24H); 2.28 (m, 1H); 2.4 (s, 3H); 2.48 (s, 3H); 2.50 (s, 3H); 6.53 (s, 1H); 6.63 (s, 1H).

45 IR (CHCl₃): 2921, 2851, 1686, 1560, 1460, 1338 cm⁻¹.

EXAMPLE 188

50

N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-2-heptylnonanoic amide

The title compound was prepared in 35% yield according to the procedure of Example 4A.

55 ¹H NMR (CDCl₃): δ 0.87 (m, 6H); 1.18-1.8 (c, 24H); 2.27 (m, 1H); 2.49 (s, 6H); 2.59 (s, 3H); 6.46 (s, 1H).

IR (CHCl₃): 2920, 2850, 1691, 1505, 1462, 1431, 1406, 1360, 1300 cm⁻¹.

EXAMPLE 1895 N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2,2-dimethyldodecanoic amide

The title compound was prepared in 49% yield according to the procedure of Example 4B.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.67 (c) and 1.32 (s) (total 24H); 2.39 (s, 3H); 2.48 (s, 3H); 2.50 (s, 3H); 6.63 (s, 1H); 6.72 (s, 1H).

10 IR (CHCl₃): 2920, 2850, 1678, 1559, 1459, 1338 cm⁻¹.

EXAMPLE 190

15

N-[2,4-bis(methylthio)pyridin-3-yl]-2,2-dimethyldodecanoic amide

The title compound was prepared in 40% yield according to the procedure of Example 4B.

20 ¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.2-1.68 (c) and 1.33 (s) (total 24H); 2.41 (s, 3H); 2.51 (s, 3H); 6.79 (s, 1H); 6.82 (d, 1H); 8.25 (d, 1H).

IR (CHCl₃): 2920, 2850, 1679, 1553, 1462, 1370 cm⁻¹.

EXAMPLE 191

25

N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-2,2-dimethyldodecanoic amide

30

The title compound was prepared in 23% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.86 (t, 3H); 1.2-1.68 (c) and 1.31 (s)(total 24H); 2.49 (s, 6H); 2.59 (s, 3H); 6.65 (s, 1H).

IR (CHCl₃): 2923, 2849, 1683, 1510, 1467, 1407, 1362, 1301 cm⁻¹.

35

EXAMPLE 192

40

N-[4,6-bis(methylthio)pyrimidin-5-yl]-2,2-dimethyldodecanoic amide

The title compound was prepared in 43% yield according to the procedure of Example 4A.

45 ¹H NMR (CDCl₃): δ 0.86 (t, 3H); 1.2-1.68 (c) and 1.32 (s)(total 24H); 2.51 (s, 6H); 6.74 (s, 1H); 8.64 (s, 1H).

IR (CHCl₃): 2924, 2851, 1688, 1522, 1468, 1406, 1359 cm⁻¹.

EXAMPLE 193

50

N-(6-methylthioquinolin-5-yl)-2,2-dimethyldodecanoic amide

55

The title compound was prepared in 4% yield according to the procedure of Example 4B.

¹H NMR (CDCl₃): δ 0.86 (t, 3H); 1.2-1.78 (c) and 1.42 (s)(total 24H); 2.55 (s, 3H); 7.44 (m, 2H); 7.66 (d, 1H); 8.07 (d, 1H); 8.13 (d, 1H); 8.83 (m, 1H).

IR (CHCl₃): 2921, 2851, 1677, 1565, 1463, 1375 cm⁻¹.

EXAMPLE 1945 N-[2,4-bis(ethylthio)-6-methylpyridin-3-yl]-tetradecanoic amide

The title compound was prepared in 68% yield according to the procedure of Example 4B.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.19-1.62 (c, 26H); 1.76 (m, 2H); 2.39 (t, 2H); 2.46 (s, 3H); 2.91 (q, 2H); 3.15 (q, 2H); 6.52 (s, 1H), 6.68 (s, 1H).

10 IR (CHCl₃): 2920, 2850, 1687, 1556, 1460 cm⁻¹.

EXAMPLE 195

15

N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-tetradecanoic amide

The title compound was prepared in 59% yield according to the procedure of Example 4B.

20 ¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.82 (c, 22H); 2.40 (s), 2.48 (s), 2.50 (s) and 2.37-2.6 (m)(total 11H); 6.50 (s, 1H); 6.64 (s, 1H).

IR (CHCl₃): 2917, 2847, 1693, 1570, 1472 cm⁻¹.

EXAMPLE 196

25

N-[4,6-bis(methylthio)pyrimidin-5-yl]-tetradecanoic amide

30

The title compound was prepared in 76% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.2-1.62 (c, 20H); 1.76 (m, 2H); 2.41 (t, 2H); 2.52 (s, 6H); 6.51 (s, 1H); 8.65 (s, 1H).

IR (CHCl₃): 2917, 2847, 1690, 1511, 1459, 1405, 1355 cm⁻¹.

35

EXAMPLE 197

40

N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]tetradecanoic amide

The title compound was prepared in 78% yield according to the procedure of Example 4A.

45 ¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.19-1.61 (c, 20H); 1.75 (m, 2H); 2.40 (t, 2H); 2.49 (s, 6H); 2.59 (s, 3H); 6.45 (s, 1H).

IR (CHCl₃): 2917, 2847, 1689, 1460, 1406, 1357 cm⁻¹.

EXAMPLE 198

50

N-(6-methylthioquinolin-5-yl)tetradecanoic amide

55

The title compound was prepared on 31% yield according to the procedure of Example 4B.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.2-1.6 (c, 20H); 1.84 (m, 2H); 2.54 (s) and 2.55 (t)(total 5H); 7.18 (s, 1H); 7.40 (m, 1H); 7.64 (d, 1H); 8.06 (m, 2H); 8.84 (b, 1H).

IR (CHCl₃): 2919, 2849, 1683, 1565, 1464, 1377 cm⁻¹.

EXAMPLE 1995 N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]pentadecanoic amide

The title compound was prepared in 53% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.81 (c, 24H); 2.4 (t, 2H); 2.5 (s, 6H); 2.6 (s, 3H); 6.44 (s, 1H).

IR (CHCl₃): 2918, 2847, 1689, 1460, 1425, 1405 cm⁻¹.

10

EXAMPLE 200

15

N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]pentadecanoic amide

The title compound was prepared in 68% yield according to the procedure of Example 4B.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.82 (c, 24H); 2.40 (s + t, 5H); 2.51 (s, 3H); 6.52 (s, 1H); 6.63 (s,

20 1H).

IR (CHCl₃): 2921, 2849, 1686, 1612, 1559, 1459 cm⁻¹.

EXAMPLE 201

25

N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]hexadecanoic amide

The title compound was prepared in 78.2% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.49 (c, 22H); 1.57 (m, 2H); 1.75 (m, 2H); 2.39 (t, 2H); 2.49 (s, 6H);

2.59 (s, 3H); 6.46 (s, 1H).

IR (CHCl₃): 2919, 2849, 1688, 1459, 1406, 1358 cm⁻¹.

35

EXAMPLE 20240 N-[4,6-bis(ethylthio)pyrimidin-5-yl]hexadecanoic amide

The title compound was prepared in 70% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.5 (c, 28H); 1.58 (m, 2H); 1.76 (m, 2H); 2.4 (t, 2H); 3.15 (q, 4H);

6.49 (s, 1H); 8.61 (s, 1H).

IR (CHCl₃): 2918, 2848, 1692, 1460, 1404, 1356 cm⁻¹.

45

EXAMPLE 203

50

N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]hexadecanoic amide

The title compound was prepared in 8.6% yield according to the procedure of Example 4.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.84 (c, 26H); 2.39 (s + t, 5H); 2.48 (s, 3H); 2.5 (s, 3H); 6.5 (s,

1H); 6.64 (s, 1H).

IR (CHCl₃): 2921, 2849, 1690, 1612, 1560, 1460 cm⁻¹.

55

EXAMPLE 2045 N-[4,6-bis (methylthio)pyrimidin-5-yl]hexadecanoic amide

The title compound was prepared in 58.8% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.49 (c, 22H); 1.57 (m, 2H); 1.76 (m, 2H); 2.41 (t, 2H); 2.51 (s, 6H); 6.54 (s, 1H); 8.65 (s, 1H).

10 IR (CHCl₃): 2920, 2849, 1696, 1521, 1465, 1407, 1358 cm⁻¹.

EXAMPLE 205

15

N-[4,6-bis(methylthio)pyrimidin-5-yl]-(Z)-9-octadecenoic amide

The title compound was prepared in 61% yield according to the procedure of Example 4A.

20 ¹H NMR (CDCl₃): δ 0.86 (t, 3H); 1.17-1.5 (c, 18H); 1.59 (m, 2H); 1.76 (m, 2H); 2.0 (c, 4H); 2.41 (t, 2H); 2.51 (s, 6H); 5.34 (m, 2H); 6.56 (s, 1H); 8.65 (s, 1H).

IR (CHCl₃): 2920, 2850, 1693, 1515, 1465, 1407, 1358 cm⁻¹.

25

EXAMPLE 206N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-(Z)-9-octadecenoic amide

30

The title compound was prepared in 55% yield according to the procedure of Example 4.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.68 (c, 20H); 1.77 (m, 2H); 2.0 (c, 4H); 2.39 (s + t, 5H); 2.47 (s, 3H); 2.49 (s, 3H); 5.34 (m, 2H); 6.51 (s, 1H); 6.63 (s, 1H).

IR (CHCl₃): 2918, 2850, 1686, 1560, 1460, 1339 cm⁻¹.

35

EXAMPLE 207

40

N-[4,6-bis(ethylthio)pyrimidin-5-yl]-(Z)-9-octadecenoic amide

The title compound was prepared in 52.3% yield according to the procedure of Example 4A.

45 ¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.19-1.5 (c, 24H); 1.58 (m, 2H); 1.76 (m, 2H); 2.01 (c, 4H); 2.40 (t, 2H); 3.15 (q, 4H); 5.34 (m, 2H); 6.5 (s, 1H); 8.61 (s, 1H).

IR (CHCl₃): 2920, 2850, 1691, 1508, 1460, 1405, 1355 cm⁻¹.

EXAMPLE 208

50

N-[2-methyl-4,6-bis(ethylthio)pyrimidin-5-yl]-(Z)-9-octadecenoic amide

55 The title compound was prepared in 66.7% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.5 (c, 24H); 1.58 (m, 2H); 1.75 (m, 2H); 2.01 (c, 4H); 2.38 (t, 2H); 2.57 (s, 3H); 3.14 (q, 4H); 5.34 (m, 2H); 6.41 (s, 1H).

IR (CHCl₃): 2919, 2849, 1690, 1459, 1407, 1357, 1312 cm⁻¹.

EXAMPLE 2095 N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-(Z)-8-octadecenoic amide

The title compound was prepared in 55% yield according to the procedure of Example 4.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.48 (c, 18H); 1.58 (m, 2H); 1.76 (m, 2H); 2.0 (c, 4H); 2.39 (t, 2H); 2.49 (s, 6H); 2.59 (s, 3H); 5.33 (m, 2H); 6.46 (s, 1H).

10 IR (CHCl₃): 2923, 2850, 1692, 1508, 1464, 1429, 1406, 1360 cm⁻¹.

EXAMPLE 210

15

N-[2,4-bis(methylthio)pyridin-3-yl]-(Z)-9-octadecenoic amide

The title compound was prepared in 43% yield according to the procedure of Example 4A.

20 ¹H NMR (CDCl₃): δ 0.86 (t, 3H); 1.18-1.5 (c, 18H); 1.59 (m, 2H); 1.77 (m, 2H); 2.01 (c, 4H); 2.41 (s + t, 5H); 2.51 (s, 3H); 5.34 (m, 2H); 6.57 (s, 1H); 6.82 (d, 1H); 8.25 (d, 1H).

IR (CHCl₃): 2920, 2850, 1687, 1552, 1461, 1375 cm⁻¹.

25

EXAMPLE 211N-[4,6-bis(methylthio)pyrimidin-5-yl]-2-dodecylthioacetamide

30

The title compound was prepared in 61% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.22-1.49 (c, 18H); 1.67 (m, 2H); 2.53 (s, 6H); 2.74 (t, 2H); 3.41 (s, 2H); 8.3 (s, 1H); 8.67 (s, 1H).

IR (CHCl₃): 2917, 2847, 1688, 1467, 1405, 1355 cm⁻¹.

35

EXAMPLE 212

40

N-[4,6-bis(ethylthio)pyrimidin-5-yl]-2-dodecylthioacetamide

The title compound was prepared in 52% yield according to the procedure of Example 4A.

45 ¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.22-1.5 (c, 24H); 1.67 (m, 2H); 2.74 (t, 2H); 3.17 (q, 4H); 3.41 (s, 2H); 8.27 (s, 1H); 8.63 (s, 1H).

IR (CHCl₃): 2918, 2848, 1687, 1466, 1404, 1353 cm⁻¹.

50

EXAMPLE 213N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-dodecylthioacetamide

55

The title compound was prepared in 45% yield according to the procedure of Example 4.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.46 (c, 18H); 1.67 (m, 2H); 2.41 (s, 3H); 2.49 (s, 3H); 2.51 (s, 3H); 2.76 (t, 2H); 3.41 (s, 2H); 6.66 (s, 1H); 8.25 (s, 1H).

IR(CHCl₃): 2918, 2848, 1678, 1561, 1476, 1337 cm⁻¹.

EXAMPLE 2145 N-[2,4-bis(methylthio)pyridin-3-yl]-2-dodecylthioacetamide

The title compound was prepared in 24% yield according to the procedure of Example 4B.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.17-1.48 (c, 18H); 1.67 (m, 2H); 2.43 (s, 3H); 2.53 (s, 3H); 2.76 (t, 2H); 3.42 (s, 2H); 6.85 (d, 1H); 8.28 (d, 1H); 8.34 (s, 1H).

10 IR (CHCl₃): 2919, 2849, 1683, 1553, 1475, 1432, 1376 cm⁻¹.

EXAMPLE 215

15

N-[2,4-bis(ethylthio)-6-methylpyridin-3-yl]-2-dodecylthioacetamide

The title compound was prepared in 37% yield according to the procedure of Example 4B.

20 ¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.47 (c, 24H); 1.67 (m, 2H); 2.47 (s, 3H); 2.77 (t, 2H); 2.92 (q, 2H); 3.15 (q, 2H); 3.41 (s, 2H); 6.69 (s, 1H); 8.24 (s, 1H).

IR (CHCl₃): 2920, 2850, 1680, 1559, 1474, 1337 cm⁻¹.

25

EXAMPLE 216N-[2,4-bis(ethylthio)pyridin-3-yl]-2-dodecylthioacetamide

30

The title compound was prepared in 27% yield according to the procedure of Example 4B.

¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.18-1.47 (c, 24H); 1.67 (m, 2H); 2.77 (t, 2H); 2.95 (q, 2H); 3.18 (q, 2H); 3.42 (s, 2H); 6.88 (d, 1H); 8.25 (d, 1H); 8.34 (s, 1H).

IR (CHCl₃): 2920, 2850, 1682, 1551, 1474, 1375 cm⁻¹.

35

EXAMPLE 217

40

N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-trans-3-nonyl-1,2,3,4-tetrahydro-2-naphthoic amide

The title compound was prepared in 7% yield according to the procedure of Example 4A. ¹H NMR (CDCl₃): δ 0.87 (m, 3H); 1.2-1.72 (c, 16H); 2.16 (m, 1H); 2.41-2.64 (c), 2.51 (s), 2.60 (s)(total 11H); 2.94-3.28 (c, 3H); 6.54 (s, 1H); 7.12 (c, 4H).

45

IR (CHCl₃): 2900, 2830, 1690, 1460 cm⁻¹.

EXAMPLE 218

50

N-[4,6-bis(methylthio)pyrimidin-5-yl]-3-nonyl-1,2,3,4-tetrahydro-2-naphthoic amide

55 The title compound was prepared in 8% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (m, 3H); 1.18-1.75 (c, 16H); 2.17 (m, 1H); 2.4-2.62 (c), 2.53 (s)(total 8H); 2.93-3.27 (c, 3H); 6.6 (s, 1H); 7.13 (c, 4H); 8.66 (s, 1H).

IR (CHCl₃): 2900, 2830, 1700, 1610, 1470 cm⁻¹.

EXAMPLE 2195 N-(2,4,6-trifluorophenyl)-trans-3-nonyl-1,2,3,4-tetrahydro-2-naphthoic amide

The title compound was prepared in 15% yield according to the procedure of Example 4A.

¹H NMR (CDCl₃): δ 0.87 (m, 3H); 1.2-1.74 (c, 16H); 2.14 (m, 1H); 2.5 (m, 2H); 2.92-3.25 (c, 3H); 6.73 (m, 3H); 7.11 (m, 2H).

10 IR (CHCl₃): 2918, 2850, 1697, 1641, 1608, 1507, 1466, 1445 cm⁻¹.

EXAMPLE 220

15

N-(6-methylthioquinolin-5-yl)-2-nonyl-1,2,3,4-tetrahydro-2-naphthoic amide

The title compound was prepared in 3% yield according to the procedure of Example 4.

20 ¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.16-2.06 (c, 17H); 2.38 (m, 1H); 2.47 (s, 3H); 2.85-3.15 (c, 3H); 3.35 (d, 1H); 7.18 (m, 5H); 7.37 (s, 1H); 7.47 (d, 1H); 7.59 (d, 1H); 7.99 (d, 1H); 8.77 (s, 1H).

EXAMPLE 221

25

N-[4,6-bis(methylthio)pyrimidin-5-yl]-2-nonyl-1,2,3,4-tetrahydro-2-naphthoic amide

The title compound was prepared in 11% yield according to the procedure of Example 4A.

30 ¹H NMR (CDCl₃): δ 0.86 (t, 3H); 1.18-1.67 (c, 15H); 1.91 (m, 2H); 2.24 (m, 1H); 2.45 (s, 6H); 2.78-2.96 (c, 2H); 3.07 (m, 1H); 3.28 (d, 1H); 6.74 (s, 1H); 7.13 (s, 4H); 8.60 (s, 1H).

IR (CHCl₃): 2921, 2849, 1681, 1518, 1454, 1406, 1357 cm⁻¹.

35

EXAMPLE 22240 N'-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-N-[4-(3-methylbutyl)benzyl]-N-cycloheptylureaA. 2,4-bis(methylthio)-6-methylpyridin-3-yl isocyanate

45 A solution of 800 mg (4 mmol) 2,4-bis(methylthio)-3-amino-6-methylpyridine and 0.4 ml (2.3 mmol) trichloromethyl chloroformate in 20 ml anhydrous dioxane was refluxed under nitrogen overnight. The reaction mixture was cooled and filtered and the filtrate was concentrated to dryness in vacuo yielding 730 mg of the title compound (81% yield) as a tan colored solid.

50

B. N'-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-N-[4-(3-methylbutyl)benzyl]-N-cycloheptylurea

A solution of 135 mg (0.6 mmol) isocyanate from Example 225A and 164 mg (0.6 mmol) N-cycloheptyl-[4-(3-methylbutyl)benzylamine in 15 ml methylene chloride was refluxed under nitrogen overnight. The reaction mixture was cooled to room temperature and concentrated in vacuo. The residue was chromatographed on 200 g silica gel, eluting with 7:3 hexane/ethyl acetate to yield 140 mg (32% yield) of the title compound as an off-white solid.

¹H NMR (CDCl₃): δ 0.92 (d, 6H); 1.39-1.74 (c, 13H); 1.98 (m, 2H); 2.37 (s, 3H); 2.44 (s, 6H); 2.59 (t, 2H);

4.37 (m, 1H); 4.50 (s, 2H); 5.47 (s, 1H); 6.57 (s, 1H); 7.18 (d, 2H); 7.32 (d, 2H).

IR (CHCl₃): 2921, 2853, 1650, 1560, 1469.

The title compounds of Examples 223-227 were prepared according to the procedure of Example 222.

5

EXAMPLE 223

10 N'-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-N-[4-(2,2-dimethylpropyl)benzyl]-N-cycloheptylurea

70% yield

¹H NMR (CDCl₃): δ 0.88 (s, 9H); 1.39-1.74 (c, 10H); 1.99 (m, 2H); 2.33 (s, 3H); 2.44 (2s, 6H); 2.48 (s, 2H); 4.38 (m, 1H); 4.52 (s, 2H); 5.46 (s, 1H); 6.57 (s, 1H); 7.13 (d, 2H); 7.31 (d, 2H).

15 IR (CHCl₃): 2922, 2853, 1651, 1561, 1470 cm⁻¹.

EXAMPLE 224

20

N'-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-N-[4-(3-methylbutyl)benzyl]-N-cycloheptylurea

72% yield.

25 ¹H NMR (CDCl₃): δ 0.92 (d, 6H); 1.40-1.75 (c, 13H); 1.98 (m, 2H); 2.44 (s, 6H); 2.55 (s, 3H); 2.60 (t, 2H); 4.37 (m, 1H); 4.51 (s, 2H); 5.37 (s, 1H); 7.20 (d, 2H); 7.31 (d, 2H).

IR (CHCl₃): 2920, 2853, 1651, 1467 cm⁻¹.

30

EXAMPLE 225

35 N'-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-N-[4-(2,2-dimethylpropyl)benzyl]-N-cycloheptylurea

70% yield.

¹H NMR (CDCl₃): δ 0.89 (s, 9H); 1.39-1.77 (c, 10H); 2.00 (m, 2H); 2.42 (s, 6H); 2.48 (s, 2H); 2.55 (s, 3H); 4.39 (m, 1H); 4.51 (s, 2H); 5.37 (s, 1H); 7.15 (d, 2H); 7.30 (d, 2H).

IR (CHCl₃): 2922, 2852, 1653, 1468, 1413 cm⁻¹.

40

EXAMPLE 226

45

N'-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-N-[4-(3-methylbutyl)benzyl]-N-heptylurea

43% yield.

50 ¹H NMR (CDCl₃): δ 0.86 (m), 0.92 (d)(total 9H); 1.20-1.72 (c, 13H); 2.47 (s, 6H); 2.57 (s), 2.60 (t)(total 5H); 3.34 (t, 2H); 4.56 (s, 2H); 5.51 (s, 1H); 7.15 (d, 2H); 7.24 (d, 2H).

IR (CHCl₃): 2920, 2850, 1659, 1468, 1415 cm⁻¹.

EXAMPLE 227

55

N'-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-N-[4-(3-methylbutyl)benzyl]-N-heptylurea

32% yield.

^1H NMR (CDCl_3): δ 0.87 (m), 0.92 (d)(total 9H); 1.19-1.72 (c, 13H); 2.37 (s, 3H); 2.46 (s, 3H); 2.48 (s, 3H); 2.59 (t, 2H); 3.34 (t, 2H); 4.58 (s, 2H); 5.61 (s, 1H); 6.61 (s, 1H); 7.17 (d, 2H); 7.27 (d, 2H).

IR (CHCl_3): 2922, 2852, 1656, 1558, 1467 cm^{-1} .

5

EXAMPLE 228

10

N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-2-decylcyclopentanecarboxamide

The title compound was prepared in 27.4% yield according to the procedure of Example 4A.

^1H NMR (CDCl_3): δ 0.87 (t, 3H); 1.2-1.82 (c, 24H); 2.27 (m, 2H); 2.49 (s, 6H); 2.59 (s, 3H); 6.6 (s, 1H).

15

IR (CHCl_3): 2921, 2851, 1681, 1450, 1407, 1360 cm^{-1} .

EXAMPLE 229

20

N-[4,6-bis(methylthio)pyrimidin-5-yl]-2-decylcyclopentanecarboxamide

The title compound was prepared in 20% yield according to the procedure of Example 4A.

25

^1H NMR (CDCl_3): δ 0.87 (t, 3H); 1.21-1.82 (c, 24H); 2.28 (m, 2H); 2.51 (s, 6H); 6.69 (s, 1H); 8.64 (s, 1H).

IR (CHCl_3): 2922, 2850, 1682, 1452, 1405, 1358 cm^{-1} .

EXAMPLE 230

30

N-[4,6-bis(methylthio)pyrimidin-5-yl]-2-decylindane-2-carboxamide

The title compound was prepared in 33.7% yield according to the procedure of Example 4A.

35

^1H NMR (CDCl_3): δ 0.86 (t, 3H); 1.14-1.88 (d, 18H); 2.49 (s, 6H); 3.04 (d, 2H); 3.58 (d, 2H); 6.63 (s, 1H);

7.2 (c, 4H); 8.63 (s, 1H).

IR (CHCl_3): 2922, 2850, 1687, 1526, 1458, 1407, 1359 cm^{-1} .

40

EXAMPLE 231

45 N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-methylthiotetradecanoic amide

The title compound was prepared in 66% yield according to the procedure of Example 4B.

^1H NMR (CDCl_3): δ 0.87 (t, 3H); 1.2-1.87 (c, 21H); 2.03 (m, 1H); 2.31 (s, 3H); 2.41 (s, 3H); 2.50 (s, 3H); 2.53 (s, 3H); 3.38 (t, 1H); 6.66 (s, 1H); 8.05 (s, 1H).

50

IR (CHCl_3): 2919, 2850, 1677, 1559, 1522, 1468, 1438 cm^{-1} .

EXAMPLE 232

55

N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-2-methylthiotetradecanoic amide

The title compound was prepared in 79% yield according to the procedure of Example 4A.
¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.21-1.86 (c, 21H); 2.04 (m, 1H), 2.29 (s, 3H); 2.52 (s, 6H); 2.62 (s, 3H); 3.37 (t, 1H); 8.0 (s, 1H).
 IR (CHCl₃): 2918, 2849, 1681, 1465, 1405 cm⁻¹.

5

EXAMPLE 233

10

N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-ethylthiotetradecanoic amide

The title compound was prepared in 51% yield according to the procedure of Example 4B.
¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.2-1.7 (c, 23H); 1.8 (m, 1H); 2.06 (m, 1H); 2.41 (s, 3H); 2.50 (s, 3H); 2.53 (s, 3H); 2.8 (q, 2H); 3.48 (t, 1H); 6.66 (s, 1H); 8.13 (s, 1H).
 IR (CHCl₃): 2920, 2850, 1675, 1560, 1466 cm⁻¹.

15

EXAMPLE 234

20

N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-2-ethylthiotetradecanoic amide

The title compound was prepared in 51% yield according to the procedure of Example 4A.
¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.2-1.88 (c, 24H); 2.03 (m, 1H); 2.52 (s, 6H); 2.62 (s, 3H); 2.79 (q, 2H); 3.48 (t, 1H); 8.08 (s, 1H).
 IR (CHCl₃): 2920, 2850, 1679, 1465, 1405 cm⁻¹.

25

30

EXAMPLE 235N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-4,5-dimethyl-trans-2-heptylcyclohex-4-enecarboxamide

The title compound was prepared in 31% yield according to the procedure of Example 4A.
¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.16-2.48 (c, 24H); 2.5 (s, 6H); 2.59 (s, 3H); 6.56 (s, 1H).
 IR (CHCl₃): 2918, 2850, 1687, 1458, 1406, 1360 cm⁻¹.

40

EXAMPLE 236

45

N-[4,6-bis(methylthio)pyrimidin-5-yl]-4,5-dimethyl-trans-2-heptylcyclohex-4-enecarboxamide

The title compound was prepared in 27% yield according to the procedure of Example 4A.
¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.17-2.49 (c, 24H); 2.52 (s, 6H); 6.64 (s, 1H); 8.65 (s, 1H).
 IR (CHCl₃): 2920, 2852, 1690, 1458, 1405, 1356 cm⁻¹.

50

EXAMPLE 237

55

N-(6-methoxyisoquinolin-5-yl)-4,5-dimethyl-trans-2-heptylcyclohex-4-enecarboxamide

The title compound was prepared in 8.5% yield according to the procedure of Example 4.

^1H NMR (CDCl_3): δ 0.86 (t, 3H); 1.14-2.5 (c, 24H); 3.99 (s, 3H); 7.17 (s, 1H); 7.39 (d, 1H); 7.52 (m, 1H); 7.94 (d, 1H); 8.45 (m, 1H); 9.14 (m, 1H).

IR (CHCl_3): 2920, 2850, 1678, 1625, 1464, 1382 cm^{-1} .

5

EXAMPLE 238

10

3-Amino-2,4-bis(methylthio)-6-methylpyridine

To a solution of 15.5 g (0.22 mol) sodium methanethiolate in 200 ml methanol was added slowly with stirring under nitrogen a solution of 20.8 g (0.1 mol) 3-nitro-2,4-dichloro-6-methylpyridine in 150 ml methanol. A precipitate formed and the mixture was stirred overnight at room temperature. The mixture was then filtered and the solid was washed first with methanol and then with water. 3-Nitro-2,4-bis(methylthio)-6-methylpyridine (18.9 g, 82% yield) was obtained as a yellow solid, mp 172-176 °C.

^1H NMR (CDCl_3): δ 2.45 (s, 3H); 2.51 (s, 3H); 2.55 (s, 3H); 6.77 (s, 1H).

A mixture of 18.9 g (0.082 mol) 3-nitro-2,4-bis(methylthio)-6-methylpyridine and 18.9 g Raney nickel in 600 ml 1,4-dioxane and 300 ml methanol was shaken with hydrogen (15 psi) in a Parr hydrogenation apparatus for 3.5 hr. The catalyst was filtered and the filtrate was concentrated to dryness in vacuo. The solid residue was chromatographed on silica gel (650g), eluting with 9:1 hexane/ethyl acetate to yield 14.0g (85% yield) of the title compound as an off-white solid.

NMR (CDCl_3): δ 2.42 (s, 3H); 2.44 (s, 3H); 2.59 (s, 3H); 4.02 (b, 2H); 6.72 (s, 1H).

The title compounds of Examples 239-241 were prepared according to the procedure of Example 238.

EXAMPLE 239

30

A. 3-Amino-2,4-bis(methylthio)pyridine

(79% yield)

^1H NMR (CDCl_3): δ 2.45 (s, 3H); 2.60 (s, 3H); 4.14 (b, 2H); 6.88 (d, 1H); 7.90 (d, 1H).

EXAMPLE 240

40

B. 3-Amino-2,4-bis(ethylthio)pyridine

(86% yield)

^1H NMR (CDCl_3): δ 1.29 (t, 3H); 1.34 (t, 3H); 2.91 (q, 3H); 3.21 (q, 3H); 4.30 (b, 2H); 6.93 (d, 1H); 7.86 (d, 1H).

EXAMPLE 241

50

C. 3-Amino-2,4-bis(ethyl)-6-methylpyridine

(86% yield)

^1H NMR (CDCl_3): δ 1.30 (t, 3H) 1.32 (t, 3H); 2.40 (s, 3H); 2.90 (q, 2H); 3.18 (q, 2H); 4.18 (b, 2H); 6.79 (s, 1H).

EXAMPLE 2425 (2S)-N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-hexylthiodecanoic amide

(S)-(-)-2-Hexylthiodecanoic, prepared according to Example 1B, was coupled with 3-amino-2,4-bis-(methylthio)-6-methylpyridine by the procedure of Example 4B to yield the title compound in 55% yield; $[\alpha]_D^{25} = -59^\circ$ (CH₃OH). A sample recrystallised from petroleum ether had mp 81-83 °C and $[\alpha]_D^{25} = -66^\circ$ (CH₃OH).
10

EXAMPLE 243

15

(2R)-N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-hexylthiodecanoic amide

The title compound was prepared in 30.1% yield according to a procedure similar to that of Example 243. A sample recrystallised from petroleum ether had mp 80-82 °C and $[\alpha]_D^{25} = +61.7^\circ$ (CH₃OH).
20

EXAMPLES 244

25

(2S)-N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide

The title compound was prepared in 47.2% yield by the coupling of S-(-)-2-hexylthiodecanoic acid with 5-amino-4,6-bis(methylthio)-2-methylpyrimidine according to the procedure of Example 4A. A sample recrystallized from diethyl ether had mp 98-100 °C and $[\alpha]_D^{25} = -62^\circ$ (CH₃OH).
30

EXAMPLE 245

35

(2R)-N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide

The title compound was prepared in 50.3% yield by a procedure similar to that of Example 245. A sample recrystallized from diethyl ether had mp 95-97.5 °C and $[\alpha]_D^{25} = +56.0^\circ$ (CH₃OH).
40

EXAMPLE 246

45

(2S)-N-[6-(methylthio)quinolin-5-yl]-2-hexylthiodecanoic amide

The title compound was prepared by the recoupling of S-(-)-2-hexylthio-decanoic acid with 5-amino-6-methylthioquinoline (Example 34) according to the procedure of Example 25. M.p.: 114-115 °C.
50

Optical rotation: $[\alpha]_D^{25} = -52^\circ$ (in CDCl₃).

Analytical analysis of enantiomeric purity was accomplished using a Chiracel OD HPLC column.

55

EXAMPLE 247

N-(6-methylthioquinolin-5-yl)-2-bromodecanamide

To a stirred solution of 2-bromodecanoic acid (184 mg, 0.73 mmol) in CH₂Cl₂ (3.0 ml) was injected by syringe oxalyl chloride (0.06 ml, 105 mol %) and then DMF (1 drop). After stirring at room temperature for 1 hour, N-methyl morpholine (0.24 ml, 300 mol %) was added. To this mixture was injected a solution of 5-amino-6-methylthioquinoline (139 mg, 100 mol%) in CH₂Cl₂ (2.0 ml). After stirring at room temperature for an additional 30 min. the reaction mixture was diluted with CH₂Cl₂ (30 ml), poured over 1.0 M H₃PO₄ (100 ml), and extracted with CH₂Cl₂ (3 x 30 ml). The combined extracts were dried (Na₂SO₄), evaporated, and chromatographed using 1% triethylamine:ethyl acetate as eluants to give the title compound (192 mg, 62% yield).

IR (CDCl₃) 3350, 2930-2820, 1710 cm⁻¹.

¹H NMR (CDCl₃) δ 8.84 (m, 1H); 8.05 (m, 3H); 7.63 (d, 1H, J=9.4 Hz); 7.41 (dd, 1H, J=3.7, 8.5 Hz); 4.57 (dd, 1H, J=5.5, 8.3 Hz); 2.54 (s, 3H); 2.25 (m, 1H); 2.15 (m, 1H); 1.58 (m, 2H); 1.26 (m, 10H); 0.85 (m, 3H).

EXAMPLE 248N-(6-methylthioquinolin-5-yl)-2-hexylaminodecanamide

A mixture of N-(6-methylthioquinolin-5-yl)-2-bromodecanamide (200 mg, 0.47 mmol) and n-hexylamine (10 ml) was heated at 120 °C for 1 hour, cooled to room temperature, and chromatographed using 1:24:25/triethylamine:ethyl acetate: hexane as eluants to give the title compound as a pink oil (184 mg, 88% yield).

IR (CHCl₃) 3280, 2940-2860, 1680 cm.

¹H NMR (CDCl₃) δ 9.38 (s, 1H); 8.80 (m, 1H); 7.97 (m, 2H); 7.62 (d, 1H, J=8.8 Hz); 7.36 (dd, 1H, J=4.3, 8.6Hz); 3.28 (dd, 1H, J=4.7, 8.6 Hz); 2.86 (m, 1H), 2.78 (m, 1H); 2.51 (s, 3H); 1.98 (m, 1H); 1.74 (m, 1H); 1.56 (m, 5H); 1.30 (m, 16H); 0.86 (m, 6H). Mass spectrum m/e (relative intensity): M⁺ 444.30 (82), 226.40 (100).

Anal. Calc'd for C₂₆H₄₁N₃OS: C, 70.4; H, 9.3; N, 9.5;

Found: C, 70.9; H, 9.4; N, 9.5.

EXAMPLE 249N-(6-methylthioquinolin-5-yl)-2-N,N-[(acyl)(hexyl)]aminodecanamide

To a stirred solution of N-(6-methylthioquinolin-5-yl)-2-hexylaminodecanamide (200 mg, 0.45 mmol) pyridine (10 ml) was added in one portion acetic anhydride (2.0 ml). After stirring at room temperature for 1 hour, the mixture was poured over 1.0 M H₃PO₄ (200 ml) and extracted with CH₂Cl₂ (3 x 100 ml). The combined extracts were dried (Na₂SO₄), evaporated, and chromatographed using ethyl acetate as eluant to give the title compound (200 mg, 91% yield).

¹H NMR (CDCl₃) δ 8.81 (d, 1H, J=2.9 Hz); 8.66 (s, 1H); 8.01 (d, 1H, J=6.7 Hz); 8.00 (d, 1H, J=2.0 Hz); 7.60 (d, 1H, J=9.1 Hz); 7.36 (dd, 1H, J=4.2, 8.6 Hz); 4.95 (m, 1H), 3.33 (t, 2H, J=8.3 Hz); 2.50(s, 3H); 2.24 (s, 3H); 2.15 (m, 1H); 1.88 (m, 1H); 1.66 (m, 2H); 1.25 (m, 18H); 0.86 (m, 6H).

Mass spectrum m/e (relative intensity): M⁺ 486.3(87), 296.3 (62), 268.3(32), 226.3 (100).

EXAMPLE 250N-(6-methylthioquinolin-5-yl)-2-N,N-[(hexyl)(methylsulfonyl)]aminodecanamide

To a stirred solution of N-(6-methylthioquinolin-5-yl)-2-hexylaminodecanamide (200 mg, 0.45 mmol) and triethylamine (0.19 ml, 300 mol%) in CH₂Cl₂ (5 ml) was added dropwise a mixture of methanesulfonyl chloride (0.038 ml, 110 mol%) in CH₂Cl₂ (5 ml) at 0 °C. After stirring at 0 °C for 30 min, the reaction mixture was allowed to warm to room temperature. After stirring at room temperature for 3 hours, the mixture was chromatographed using 1% triethylamine:ethyl acetate as eluants to give the title compound (120 mg, 51% yield).

¹H NMR (CDCl₃) δ 8.84 (s, 1H); 8.31 (s, 1H); 8.13 (d, 1H, J=8.4 Hz); 8.05 (d, 1H, J=9.0 Hz); 7.64 (d, 1H, J=9.0 Hz); 7.42 (dd, 1H, J=4.2, 8.5 Hz); 4.51 (t, 1H, J=7.4 Hz); 3.40 (m, 3H); 3.01 (s, 3H); 2.54 (s, 3H); 2.19 (m, 1H); 1.77 (m, 2H); 1.28 (m, 18H); 0.87 (m, 6H).

Mass spectrum m/e (relative intensity): M⁺ 522.3 (100).

EXAMPLE 251

15

N-(6-methylthioquinolin-5-yl)-2-N,N-[(benzenesulfonyl)(hexyl)]aminodecanamide

The title compound was prepared according to the procedure of Example 250.

¹H NMR (CDCl₃) δ 8.85 (s, 1H); 8.50 (s, 1H); 8.22 (d, 1H, J=8.4 Hz); 8.11 (d, 1H, J=9.0 Hz); 7.91 (m, 2H); 7.68 (d, 1H, J=9.0 Hz); 7.59 (m, 3H); 7.45 (dd, 1H, J=4.1, 8.5 Hz); 4.45 (t, 1H, J=7.3 Hz); 3.49 (m, 2H); 3.28 (m, 1H); 2.56 (s, 3H); 2.06 (m, 1H); 1.76 (m, 2H); 1.20 (m, 18H); 0.86 (m, 6H).

Mass spectrum m/e (relative intensity): M⁺ 584.4 (100).

25

EXAMPLE 252

N-[4-dimethylamino-6-(cyano)(hexyl)aminopyrimidin-5-yl]-2-N,N-[(cyano)(hexyl)]aminodecanamide

To a stirred solution of N-[4-dimethylamino-6-(cyano)(hexyl)aminopyrimidin-5-yl]-2-hexylaminodecanamide (200 mg, 0.41 mmol), N-methyl morpholine (5 drops) in THF (5 ml) was added in one portion solid cyanobromide (95 mg, 220 mol%). After stirring at room temperature for 1 hour the mixture was chromatographed using 1:1 ethyl acetate:hexane as eluants to give the title compound (130 mg, 59% yield).

IR (CHCl₃) 3450, 3000-2800, 2200, 1740, 1650, 1600 cm⁻¹.

¹H NMR (CDCl₃) δ 8.24 (s, 1H); 4.22 (m, 1H); 4.07 (m, 2H); 3.40 (m, 2H); 3.18 (m, 1H); 2.97 (s, 3H); 2.96 (s, 3H); 2.12 (m, 1H); 1.94 (m, 1H); 1.64 (m, 2H); 1.49 (m, 2H); 1.28 (m, 24H); 0.86 (m, 8H).

Mass spectrum m/e (relative intensity): M⁺ 542 (100);

Anal. calc'd for C₃₀H₅₂N₈O: C, 66.6; H, 9.7; N, 20.7;

Found: C, 66.4; H, 9.8; N, 20.6.

EXAMPLE 253

45

N-(6-methylthioisoquinolin-5-yl)-2-(hexylthio)decanamide

Commercially available 4-chlorobenzaldehyde (10 g) was cyclized with aminoacetaldehyde diethyl acetal according to the procedure of Hendrickson, et. al., J. Org. Chem., 48, 3344 (1983), to give 6-chloroisoquinoline (600 mg). Nitration of the obtained product using the procedure of Campbell et. al., J. Am. Chem. Soc., 68, 1559 (1946) gave 6-chloro-5-nitroisoquinoline (700 mg). Substitution of 6-chloro for thiomethoxide according to the procedure of Massie, Iowa State Coll. J. Sci., 21, 41 (1946) (Ca 41:3033) gave 6-methylthio-5-nitroisoquinoline (632 mg). This material (200 mg) was reduced using stannous chloride (2.0 g) and concentrated hydrochloric acid (30 ml) to give 5-amino-6-(methylthio)isoquinoline (143 mg). This material (140 mg) was coupled with 2-hexylthiodecanoic acid (prepared according to the procedures described in Examples 1 and 3) using the procedure described in Example 25 to give the title

compound (80 mg).

¹H NMR (CDCl₃) δ 9.16 (s, 1H); 8.59 (s, 1H); 8.48 (d, 1H, J=5.9 Hz); 7.88 (d, 1H, J=8.7 Hz); 7.48 (m, 2H); 3.54 (dd, 1H, J=6.3, 8.1 Hz); 2.79 (t, 2H, J=7.4Hz); 2.55 (s, 3H); 2.12 (m, 1H); 1.88 (m, 1H); 1.65 (m, 4H); 1.30 (m, 16 H); 0.88 (m, 6H).

- 5 Mass spectrum m/e (relative intensity): M⁺ 461.3;
 Anal. calc'd for C₂₆H₄₀N₂OS₂: C, 67.8; H, 8.8; N, 6.1;
 Found: C, 67.9; H, 8.9; N, 6.1.

10 EXAMPLE 254

N-[4,6-bis(dimethylamino)pyrimidin-5-yl]-cis-9-octadecenamide

- 15 5-Amino-4,6-bis(dimethylamino)pyrimidine (prepared by reacting commercially available 4,6-dichloro-5-nitro pyrimidine with excess dimethylamine followed by reduction of the nitro group according to the procedure of Jacobs et. al. , J. Am. Chem. Soc. , 42 , 2278 (1920)) was coupled with oleoyl chloride using the procedure described in Example 4 to give the title compound.

- 20 ¹H NMR (CDCl₃) δ 8.15 (s, 1H); 7.52 (s, 1H); 5.31 (m, 2H); 3.06 (s, 6H); 2.99 (s, 6H); 1.98 (m, 4H); 1.72 (m, 2H); 1.24 (m, 22H); 0.85 (m, 3H).
 Mass Spectrum m/e (relative intensity): M⁺ 446 (100), 182 (47).

25 EXAMPLE 255

N-(4-dimethylamino-6-chloropyrimidin-5-yl)-cis-9-octadecenamide

- 30 5-Amino-4-(dimethyl)amino-6-chloropyrimidine (prepared by reacting commercially available 5-amino-4,6-dichloropyrimidine with excess dimethylamine) was coupled with oleoyl chloride according to the procedure described in Example 4 to give the title compound.

- IR (CHCl₃) 3400, 3000-2800, 1700, 1570 cm⁻¹.
 35 ¹H NMR (CDCl₃) δ 8.23 (s, 1H); 6.82 (s, 1H); 5.34 (m, 2H); 3.15 (s, 6H); 2.40 (t, 2H, J = 7.6 Hz); 2.01 (m, 4H); 1.73 (m, 2H); 1.25 (m, 20H); 0.86 (m, 3H).
 Mass spectrum m/e (relative intensity): M⁺ 437 (100). Anal. Calc'd for C₂₄H₄₁N₄OCl: C, 66.0; H, 9.5; N, 12.8. Found: C, 65.7; H, 9.5; N, 12.9.

40 EXAMPLE 256

N-(4-dimethylamino-6-methylthiopyrimidin-5-yl)-cis-9-octadecenamide

- 45 5-Amino-4-dimethylamino-6-methylthiopyrimidine (prepared by reacting 5-amino-4-dimethylamino-6-chloropyrimidine with sodium thiomethoxide) was coupled with oleoyl chloride using the procedure described in Example 4 to give the title compound.

- 50 IR (KBr) 3220, 3000-2800, 1650, 1550 cm⁻¹.
¹H NMR (CDCl₃) δ 8.32 (s, 1H); 6.68 (s, 1H); 5.33 (m, 2H); 3.09 (s, 6H); 2.45 (s, 3H); 2.37 (t, 2H, J = 7.7 Hz); 2.00 (m, 4H); 1.72 (m, 2H); 1.25 (m, 20H); 0.86 (m, 3H).
 Mass spectrum m/e (relative intensity): M⁺ 449.4 (62), 185 (100). Anal. Calc'd for C₂₅H₄₄N₄OS: C, 66.9; H, 9.9; N, 12.5. Found: C, 67.3; H, 10.2; N, 12.4.

55 EXAMPLE 257

N-(4-bromoisoquinolin-5-yl)-2-hexylthiodecanamide

5-Amino-4-bromoisoquinoline (prepared according to the process described by Gordon et al., J. Het. Chem., 4, 410 (1967)) was coupled with 2-hexylthiodecanoic acid (prepared by the procedures described in Examples 1 and 3) using the procedure described in Example 25 to give the title compound.

IR (CHCl₃) 3320, 3000-2800, 1680 cm⁻¹.

¹H NMR (CDCl₃) δ 10.10 (s, 1H); 9.11 (s, 1H); 8.68 (s, 1H); 8.51 (d, 1H, J = 7.7 Hz); 7.82 (dd, 1H, J = 1.2, 8.1 Hz); 7.67 (t, 1H, J = 8.0 Hz); 3.50 (t, 1H, J = 7.3 Hz); 2.60 (t, 2H, J = 7.3 Hz); 2.05 (m, 1H); 1.82 (m, 1H); 1.59 (m, 4H); 1.24 (m, 16H); 0.85 (m, 6H).

Mass spectrum m/e (relative intensity): M⁺ 495 (50), 415 (100). Anal. calc'd for C₂₅H₃₇N₂OSBr: C, 60.8; H, 7.6; N, 5.7. Found: C, 61.0; H, 7.5; N, 5.7.

EXAMPLE 258N-(4-methylthioisoquinolin-5-yl)-2-hexylthiodecanamide

5-Amino-4-bromoisoquinoline (prepared according to the process described by Gordon et al., J. Het. Chem., 4, 410 (1967)) was allowed to react with sodium thiomethoxide according to the procedure of Massie, Iowa State Coll. J. Sci., 21, 41 (1946) (Ca. 41: 3044 g) to give 5-amino-4-methylthioisoquinoline. This material was coupled with 2-hexylthiodecanoic acid (prepared by the procedures described in Examples 1 and 3) using the procedure described in Example 25 to give the title compound.

IR (KBr) 3230, 3000-2800, 1660 cm⁻¹.

¹H NMR (CDCl₃) δ 11.65 (s, 1H); 9.12 (s, 1H); 8.84 (d, 1H, J = 7.8 Hz); 8.63 (s, 1H); 7.76 (dd, 1H, J = 1.2, 8.1 Hz); 7.64 (t, 1H, J = 8.0 Hz); 3.44 (t, 1H, J = 7.5 Hz); 2.65 (m, 2H); 2.53 (s, 3H); 2.03 (m, 1H); 1.83 (m, 1H); 1.59 (m, 4H); 1.24 (m, 16H); 0.84 (m, 6H).

Mass spectrum m/e (relative intensity): M⁺ 461.3 (100). Anal. calc'd for C₂₆H₄₀N₂OS₂: C, 67.8; H, 8.8; N, 6.1. Found: C, 68.0; H, 8.9; N, 6.0.

EXAMPLE 259N-(4-bromoisoquinolin-5-yl)-cis-9-octadecenamide

5-Amino-4-bromoisoquinoline (prepared according to the process described by Gordon et al., J. Het. Chem., 4, 410 (1967)) was coupled with oleoyl chloride using the procedure described in Example 4, to give the title compound.

IR (CHCl₃) 3420, 3000-2800, 1695 cm⁻¹.

¹H NMR (CDCl₃) δ 9.10 (s, 1H); 9.04 (s, 1H); 8.65 (s, 1H); 8.42 (d, 1H, J = 7.3 Hz); 7.80 (d, 1H, J = 8.0 Hz); 7.65 (t, 1H, J = 7.9 Hz); 5.33 (m, 2H); 2.49 (t, 2H, J = 7.3 Hz); 2.00 (m, 3H); 1.80 (m, 2H); 1.25 (m, 21H); 0.86 (m, 3H).

Mass spectrum m/e (relative intensity): M⁺ 489.3 (37), 461.3 (14), 407.3 (100). Anal. calc'd for C₂₇H₃₉N₂OBr: C, 66.8; H, 8.1; N, 5.8. Found: C, 67.3; H, 8.4; N, 5.6.

EXAMPLE 260N-(4-dimethylamino-6-hexylaminopyrimidin-5-yl)-2-hexylaminodecanamide

5-Amino-4-(dimethyl)amino-6-chloropyrimidine (prepared as described in Example 257) was coupled with 2-bromodecanoic acid using the procedure described in Example 249, followed by the addition of n-hexylamine as described in Example 250 to give the title compound.


IR (CHCl₃) 3280, 3000-2700, 1660, 1600 cm⁻¹.

¹H NMR (CDCl₃) δ 8.86 (s, 1H); 8.20 (s, 1H); 5.47 (t, 1H, J = 5.0 Hz); 3.40 (m, 2H); 3.18 (m, 1H); 2.88 (s, 6H); 2.66 (m, 2H); 1.85 (m, 1H); 1.44 (m, 31H); 0.86 (m, 8H).

Mass spectrum m/e (relative intensity): M⁺ 491.5 (100). Anal. calc'd for C₂₈H₅₄N₆O: C, 68.5; H, 11.1; N, 17.1. Found: C, 68.8; H, 11.3; N, 17.5.





Claims

1. A compound of the formula



wherein Q is -CR²R³R⁴ or -NR¹⁷R¹⁸:

R¹ is



R², R³ and R⁴ may be the same or different, and

(a) are selected from the group consisting of hydrogen, (C₁-C₄) alkyl, A, XR¹⁰, phenyl-(C₁-C₇) alkyl, and (C₅-C₆) cycloalkyl-(C₁-C₆) alkyl, with the proviso that at least one of R², R³ and R⁴ must be A, and with the proviso that when R¹ is a group of the formula XXVII or a group of the formula XXVI wherein G is nitrogen and wherein neither R⁵, R⁶ nor R¹⁵ is NR¹⁹R²⁰, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄) alkylthio, phenylthio or heteroalkylthio, either at least one of R², R³ and R⁴ must be XR¹⁰, or two of R², R³ and R⁴ must be A; or

(b) R² and R³ together with the carbon to which they are attached form a cyclic or bicyclic system selected from the group consisting of (C₃-C₇) cycloalkyl, (C₃-C₇) cycloalkenyl, (C₆-C₁₄) bicycloalkyl, (C₆-C₁₄) bicycloalkenyl, and aryl-fused and heteroaryl-fused systems containing 8 to 15 carbon atoms, one ring of any of said aryl-fused and heteroaryl-fused systems being aromatic and the ring containing the carbon to

which R² and R³ are attached being non-aromatic, one of the carbons of said aromatic ring being optionally replaced by sulfur or oxygen, one or more carbons of said non-aromatic ring being optionally replaced by sulfur or oxygen, and one or more carbons of said aromatic ring being optionally replaced by nitrogen; one or two carbons of said cycloalkyl or bicycloalkyl groups being optionally replaced by sulfur or oxygen, and said cyclic or bicyclic system being optionally substituted with one to five substituents independently selected from the group consisting of phenyl, substituted phenyl, (C₁-C₆) alkyl and A, with the proviso that one and only one of said substituents is A, and one and only one of said substituents is phenyl or substituted phenyl, said substituted phenyl being substituted with one or more substituents independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) alkylthio, halogen and trifluoromethyl; and R⁴ is hydrogen, XR¹⁰ or A; with the proviso that when R¹ is a group of the formula XXVII or a group of the formula XXVI wherein G is nitrogen and wherein neither R⁵, R⁶ nor R¹⁵ is NR¹⁹R²⁰, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄) alkylthio, phenylthio or heteroalkylthio, R² and R³, together with the carbon to which they are attached, do not form a (C₃-C₇) cycloalkyl ring containing only carbon atoms:

A is a hydrocarbon containing 4 to 16 carbons and 0, 1 or 2 double bonds:

X is O, S, SO, SO₂, NH, NR²³CO or NSO₂R²⁴, wherein R²³ is hydrogen or (C₁-C₆) alkyl and R²⁴ is (C₁-C₆) alkyl, phenyl or (C₁-C₃) alkyl-phenyl:

R⁵, R⁶, R¹⁵ and R¹⁶ are each independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, iodo, (C₁-C₄) alkyl, (C₁-C₄) haloalkyl, (C₁-C₄) alkoxy, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄) alkylthio, substituted phenylthio, heteroarylthio, heteroaryloxy, and NR¹⁹R²⁰ wherein R¹⁹ and R²⁰ are the same or different and are selected from the group consisting of hydrogen, (C₁-C₄) alkyl, phenyl, substituted phenyl, (C₁-C₄) acyl, aroyl, and substituted aroyl, wherein said substituted phenyl and substituted aroyl groups are substituted with one or more substituents independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) alkylthio, halogen and trifluoromethyl, or R¹⁹ and R²⁰, together with the nitrogen to which they are attached, form a piperidine or morpholine ring; and wherein R⁵, R⁶, R¹⁵ and R¹⁶, when attached to a bicyclic system, may be attached to either ring of such system, with the proviso that no more than 3 non-hydrogen substituents may be attached to any one ring of such system:

R⁷, R⁸ and R⁹ are the same or different; R⁷ is selected from the group consisting of hydrogen, (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, methyl and fluoro; and R⁸ and R⁹ are each independently selected from the group consisting of (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, methyl, and fluoro:

R¹⁰ is selected from the group consisting of (C₄-C₁₂) cycloalkyl, (C₄-C₁₂) straight or branched alkyl, (C₄-C₁₂) cycloalkyl-(C₁-C₆) alkyl, phenyl-(C₁-C₆) alkyl, (substituted phenyl)-(C₁-C₆) alkyl, (C₁-C₆) alkyl-phenyl, (C₁-C₆) alkyl-(substituted phenyl), substituted thiazoles, substituted benzothiazoles, and substituted pyridines; wherein the substituents on the substituted phenyl, substituted thiazoles, substituted benzothiazoles and substituted pyridines are selected from the group consisting of (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, (C₁-C₆) alkyl, halo and trifluoromethyl:


B, D, E and G are selected from the group consisting of nitrogen and carbon, with the proviso that one or more of B, D and E is nitrogen, and with the proviso that when G is nitrogen, the group XXVI is attached to the nitrogen of formula I at the 4 or 5 position of the pyrimidine ring (designated by a and b):

and R¹⁷ and R¹⁸ are each independently selected from the group consisting of (C₄-C₁₂) straight or branched alkyl, phenyl-(C₁-C₄) alkyl, and (C₁-C₆) alkylphenyl-(C₁-C₆) alkyl:

with the proviso that when Q is NR¹⁷R¹⁸, R¹ is a group of the formula XXVI or XXIV, or a group of the formula XXVII wherein R⁷, R⁸ and R⁹ are each methoxy:

or pharmaceutically acceptable salt of said compound.

2. A compound according to claim 1, wherein Q is -CR²R³R⁴ and R¹ is




3. A compound according to claim 1, said compound being selected from the group consisting of:

N-(2,4,6-trifluorophenyl)-2-(hexylthio)octanoic amide;


N-(2,4,6-trimethoxyphenyl)-2-(hexylthio)octanoic amide;

- N-(2,4,6-trimethoxyphenyl)-2-(6-ethoxybenzothiazol-2-yl)decanoic amide;
 \bar{N} -(6-methoxyquinolin-5-yl)-2-(hexylthio)decanoic amide;
 \bar{N} -(6-methylthioquinolin-5-yl)-2-(hexylthio)decanoic amide;
 \bar{N} -(6-methylthio-8-acetaminoquinolin-5-yl)-2-(hexylthio)decanoic amide;
 5 \bar{N} -(6-methoxyisoquinolin-5-yl)-2-(hexylthio)decanoic amide;
 \bar{N} -(6-methylthioisoquinolin-5-yl)-2-(hexylthio)decanoic amide;
 \bar{N} -(6-methylisoquinolin-5-yl)-2-(hexylthio)decanoic amide;
 \bar{N} -(6-methylthioquinolin-5-yl)-2-(4-(3-methylpropyl)phenoxy)nonanoic amide;
 $(2S)$ -N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-hexylthiodecanoic amide;
 10 $(2S)$ -N-[2-methyl-4,6-bis(methylthio)pyrimidin-5-yl]-2-hexylthiodecanoic amide;
 $(2S)$ -N-[6-(methylthio)quinolin-5-yl]-2-hexylthiodecanoic amide;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-2-hexylthiodecanoic amide;
N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2-hexylthiodecanoic amide;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-2,2-dimethyldodecanoic amide;
 15 N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-2,2-dimethyldodecanoic amide;
N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-N-[4-(3-methylbutyl)benzyl]-N-cycloheptylurea;
N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-N-[4-(3-methylbutyl)benzyl]-N-heptylurea;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-N-[4-(3-methylbutyl)benzyl]-N-cycloheptylurea;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-N-[4-(3-methylbutyl)benzyl]-N-heptylurea;
 20 N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-4,5-dimethyl-trans-2-heptylcyclohex-4-ene-carboxamide;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-2-heptylnonanoic amide;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-pentadecanoic amide;
N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-pentadecanoic amide;
N-[2,4-bis(methylthio)-6-methylpyridin-3-yl]-(Z)-9-octadecenoic amide;
 25 N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-(Z)-9-octadecenoic amide;
N-[4,6-bis(methylthio)-2-methylpyrimidin-5-yl]-trans-3-nonyl-1,2,3,4-tetrahydro-2-naphthoic amide;
N-[4,6-bis(methylthio)pyrimidin-5-yl]-trans-3-nonyl-1,2,3,4-tetrahydro-2-naphthoic amide;
N-(2,4,6-trimethoxyphenyl)-2-nonyl-1,2,3,4-tetrahydronaphth-2-yl carboxamide;
 \bar{N} -(2,4,6-trifluorophenyl)-2-nonyl-1,2,3,4-tetrahydronaphth-2-yl carboxamide;
 30 \bar{N} -(2,4,6-trifluorophenyl)-2-nonylindan-2-yl carboxamide;
 \bar{N} -(2,4,6-trimethoxyphenyl)-6-heptyl-3,4-dimethylcyclohex-2-enyl carboxamide;
 \bar{N} -(2,4,6-trimethoxyphenyl)-2-nonylbicyclo[2.2.1]hept-5-en-2-yl carboxamide;
 \bar{N} -(6-methylthioquinolin-5-yl)-2-octyl-1,3-dithian-2-yl carboxamide;
 \bar{N} -(2,4,6-trimethoxyphenyl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea
 35 \bar{N} -(2,4,6-trimethoxyphenyl)- \bar{N} -(4-(2,2-dimethylpropyl)phenylmethyl)-N'-heptylurea;
 \bar{N} -(isoquinolin-5-yl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea;
 \bar{N} -(quinolin-5-yl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea; and
 \bar{N} -(6-methoxyquinolin-5-yl)-N'-(4-(3-methylbutyl)phenylmethyl)-N'-heptylurea.
 4. A compound of the formula



wherein

- Z is hydroxy, chloro or bromo; R^2 , R^3 and R^4 are each independently selected from the group consisting of
 (C₆-C₁₆) alkyl, (C₆-C₁₆) alkenyl containing 1-2 double bonds, phenyl (C₁-C₇) alkyl, (C₅-C₆) cycloalkyl (C₁-
 50 (C₆-C₁₆) alkyl, XR¹⁰ and hydrogen, with the proviso that at least one of R^2 , R^3 and R^4 is (C₆-C₁₆) alkyl or
 (C₆-C₁₆) alkenyl containing 1-2 double bonds; or
 R^2 and R^3 together with the carbon to which they are attached form a cyclic or bicyclic ring system
 selected from the group consisting of



and aryl-fused and heteroaryl fused systems containing 8 to 15 carbon atoms, one ring of any of said aryl-fused and heteroaryl-fused systems being aromatic and the other ring containing the carbon to which R^2 and R^3 are attached being non-aromatic, one of the carbons of said aromatic ring being optionally replaced by sulfur or oxygen, one or more carbons of said non-aromatic ring being optionally replaced by sulfur or oxygen, and one or more carbons of said aromatic ring being optionally replaced by nitrogen; and wherein R^4 is hydrogen, A or XR^{10} ; A is a hydrocarbon containing 4 to 16 carbons and 0, 1 or 2 double bonds; X is oxygen or sulfur; R^{10} is (C_4-C_{12}) straight or branched alkyl, phenyl substituted with (C_1-C_6) alkyl or phenyl (C_1-C_4) alkyl; R^{11} is (C_6-C_{12}) alkyl or (C_6-C_{12}) alkenyl containing 1-2 double bonds; n and m are each independently 1 or 2; o and p are each independently 0, 1 or 2; each dotted line represents an optional double bond; and each asterisk represents the carbon to which R^2 and R^3 are attached; with the proviso that when R^2 and R^3 form any of ring systems XVIII, XIX and XX, R^4 is A, and when R^2 and R^3 form any of ring systems XXI, XXII and XXIII, R^4 is hydrogen; and with the proviso that when R^2 and R^3 do not form a ring system, no more than one of R^2 , R^3 and R^4 is hydrogen.

5. A compound according to claim 4 selected from the group consisting of:

- 2-(hexylthio)octanoic acid;
- 2-(hexylthio)nonanoic acid;
- 2-(hexylthio)decanoic acid;
- 2-(heptylthio)octanoic acid;
- 2-(heptylthio)nonanoic acid;
- 2-(heptylthio)decanoic acid;
- 2-nonylindan-2-yl carboxylic acid;
- 2-decylindan-2-yl carboxylic acid;
- 2-octyl-1,2,3,4-tetrahydronaphth-2-yl carboxylic acid; and
- 2-nonyl-1,2,3,4-tetrahydronaphth-2-yl carboxylic acid;

6. A pharmaceutical composition for inhibiting acyl coenzyme A: cholesterol acyltransferase, inhibiting intestinal absorption of cholesterol, reversing or slowing the development of atherosclerosis, or lowering the concentration of serum cholesterol in a mammal, comprising an amount of a compound according to claim 1 that is effective in inhibiting acyl coenzyme A: cholesterol acyltransferase or intestinal absorption of


cholesterol, or is effective in reversing or slowing the development of atherosclerosis or lowering the concentration of serum cholesterol, and a pharmaceutically acceptable carrier.

7. A method for inhibiting acyl coenzyme A:


cholesterol acyltransferase, inhibiting intestinal absorption of cholesterol, reversing or slowing the development of atherosclerosis, or lowering the concentration of serum cholesterol in a mammal, comprising an amount of a compound according to claim 1 that is effective in inhibiting acyl coenzyme A:

cholesterol acyltransferase or intestinal absorption of cholesterol, or is effective in reversing or slowing the development of atherosclerosis or lowering the concentration of serum cholesterol and a pharmaceutically acceptable carrier.

8. A compound according to claim 1, wherein Q is $-CR^2R^3R^4$ and R^1 is




9. A compound of the formula




I

wherein Q is $-CR^2R^3R^4$ or $-NR^{17}R^{18}$:


R^1 is



XXIV




XXV



XXVI

or



XXVII

R², R³ and R⁴ may be the same or different, and

(a) are selected from the group consisting of hydrogen, (C₁-C₄) alkyl, A, XR¹⁰, phenyl-(C₁-C₇) alkyl, and (C₅-C₆) cycloalkyl-(C₁-C₆) alkyl, with the proviso that at least one of R², R³ and R⁴ must be A, and with the proviso that when R¹ is a group of the formula XXVII or a group of the formula XXVI wherein G is nitrogen and wherein neither R⁵, R⁶ nor R¹⁵ is NR¹⁹R²⁰, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl(C₁-C₄) alkylthio, phenylthio or heteroalkylthio, either at least one of R², R³ and R⁴ must be XR¹⁰, or two of R², R³ and R⁴ must be A; or

(b) R² and R³, together with the carbon to which they are attached form a cyclic or bicyclic system selected from the group consisting of (C₃-C₇) cycloalkyl, (C₃-C₇) cycloalkenyl, (C₆-C₁₄) bicycloalkyl, (C₆-C₁₄) bicycloalkenyl, and aryl-fused and heteroaryl-fused systems containing 8 to 15 carbon atoms, one ring of any of said aryl-fused and heteroaryl-fused systems being aromatic and the ring containing the carbon to which R² and R³ are attached being non-aromatic, one of the carbons of said aromatic ring being optionally replaced by sulfur or oxygen, one or more carbons of said non-aromatic ring being optionally replaced by sulfur or oxygen, and one or more carbons of said aromatic ring being optionally replaced by nitrogen; one or two carbons of said cycloalkyl or bicycloalkyl groups being optionally replaced by sulfur or oxygen, and said cyclic or bicyclic system being optionally substituted with one to five substituents independently selected from the group consisting of phenyl, substituted phenyl, (C₁-C₆) alkyl and A, with the proviso that one and only one of said substituents is A, and one and only one of said substituents is phenyl or substituted phenyl, said substituted phenyl being substituted with one or more substituents independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) alkylthio, halogen and trifluoromethyl; and R⁴ is hydrogen, XR¹⁰ or A; with the proviso that when R¹ is a group of the formula XXVI or XXVII, R² and R³, together with the carbon to which they are attached, do not form a (C₃-C₇) cycloalkyl or (C₆-C₇) bicycloalkyl ring containing only carbon atoms:

A is a hydrocarbon containing 4 to 16 carbons and 0, 1 or 2 double bonds:

X is O, S, SO, SO₂, NH, NR²³CO or NSO₂R²⁴, wherein R²³ is hydrogen or (C₁-C₆) alkyl and R²⁴ is (C₁-C₆) alkyl, phenyl or (C₁-C₃) alkyl-phenyl:

R⁵, R⁶, R¹⁵ and R¹⁶ are each independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, iodo, (C₁-C₄) alkyl, (C₁-C₄) haloalkyl, (C₁-C₄) alkoxy, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄) alkylthio, substituted phenylthio, heteroarylthio, heteroaryloxy, and NR¹⁹R²⁰, wherein R¹⁹ and R²⁰ are the same or different and are selected from the group consisting of hydrogen, (C₁-C₄) alkyl, phenyl, substituted phenyl, (C₁-C₄) acyl, aroyl, and substituted aroyl, wherein said substituted phenyl and substituted aroyl groups are substituted with one or more substituents independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) alkylthio, halogen and trifluoromethyl or R¹⁹ and R²⁰, together with the nitrogen to which they are attached, form a piperidine or morpholine ring; and wherein R⁵, R⁶, R¹⁵ and R¹⁶, when attached to a bicyclic system, may be attached to either ring of such system, with the proviso that no more than 3 non-hydrogen substituents may be attached to any one ring of such system:

R⁷, R⁸ and R⁹ are the same or different; R⁷ is selected from the group consisting of hydrogen, (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, methyl and fluoro; and R⁸ and R⁹ are each independently selected from the group consisting of (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, methyl, and fluoro:

R¹⁰ is selected from the group consisting of (C₄-C₁₂) cycloalkyl, (C₄-C₁₂) straight or branched alkyl, (C₄-C₁₂) cycloalkyl-(C₁-C₆) alkyl, phenyl-(C₁-C₆) alkyl, (substituted phenyl)-(C₁-C₆) alkyl, (C₁-C₆) alkyl-phenyl, (C₁-C₆) alkyl-(substituted phenyl), substituted thiazoles, substituted benzothiazoles, and substituted pyridines; wherein the substituents on the substituted phenyl, substituted thiazoles, substituted benzothiazoles and substituted pyridines are selected from the group consisting of (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, (C₁-C₆) alkyl, halo and trifluoromethyl:

B, D, E and G are selected from the group consisting of nitrogen and carbon, with the proviso that one or more of B, D and E is nitrogen, and with the proviso that when G is nitrogen, the group XXVI is attached to the nitrogen of formula I at the 4 or 5 position of the pyrimidine ring (designated by a and b):


and R¹⁷ and R¹⁸ are each independently selected from the group consisting of (C₄-C₁₂) straight or branched alkyl, phenyl-(C₁-C₄) alkyl, and (C₁-C₆) alkyl/phenyl-(C₁-C₆) alkyl:

with the proviso that when Q is NR¹⁷R¹⁸, R¹ is a group of the formula XXVI or XXIV, or a group of the formula XXVII wherein R⁷, R⁸ and R⁹ are each methoxy:

or pharmaceutically acceptable salt of said compound.


10. A compound according to any one of claims 2, 4 and 8, wherein:

(a) R⁴ is A, and R² and R³ together with the carbon to which they are attached form a cyclic or bicyclic ring system selected from the group consisting of




wherein n and m are each independently 1 or 2, and the asterisk designates the carbon to which R^2 and R^3 are attached; or

(b) R^4 is A, and R^2 and R^3 together with the carbon to which they are attached form the bicyclic ring system




wherein o is 0, 1 or 2, and the asterisk designates the carbon to which R^2 and R^3 are attached; or

(c) R^4 is hydrogen, and R^2 and R^3 together with the carbon to which they are attached form a cyclic or bicyclic ring system selected from the group consisting of



wherein p is 0, 1 or 2, R^{11} is A, the dotted line represents an optional double bond, and the asterisk designates the carbon to which R^2 and R^3 are attached.


11. A compound of the formula




wherein R^{21} is (C_1-C_3) alkyl and R^{22} is hydrogen or (C_1-C_3) alkyl.

Claims for the following Contracting States: ES, GR


1. A process for making compound of the formula




wherein R¹ is



XXIV




XXV



XXVI

or



XXVII

R², R³ and R⁴ may be the same or different, and

(a) are selected from the group consisting of hydrogen, (C₁-C₄) alkyl, A, XR¹⁰, phenyl-(C₁-C₇) alkyl, and (C₅-C₆) cycloalkyl-(C₁-C₆) alkyl, with the proviso that at least one of R², R³ and R⁴ must be A, and with the proviso that when R¹ is a group of the formula XXVII or a group of the formula XXVI wherein G is nitrogen and wherein neither R⁵, R⁶ nor R¹⁵ is NR¹⁹R²⁰, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄) alkylthio, phenylthio or heteroalkylthio, either at least one of R², R³ and R⁴ must be XR¹⁰, or two of R², R³ and R⁴ must be A; or

(b) R² and R³ together with the carbon to which they are attached form a cyclic or bicyclic system selected from the group consisting of (C₃-C₇) cycloalkyl, (C₃-C₇) cycloalkenyl, (C₆-C₁₄) bicycloalkyl, (C₆-C₁₄) bicycloalkenyl, and aryl-fused and heteroaryl-fused systems containing 8 to 15 carbon atoms, one ring of said aryl-fused and heteroaryl-fused systems being aromatic and the ring containing the carbon to which R² and R³ are attached being non-aromatic, one of the carbons of said aromatic ring being optionally replaced by sulfur or oxygen, one or more carbons of said non-aromatic ring being optionally replaced by sulfur or oxygen, and one or more carbons of said aromatic ring being optionally replaced by nitrogen; one or two carbons of said cycloalkyl or bicycloalkyl groups being optionally replaced by sulfur or oxygen, and said cyclic or bicyclic system being optionally substituted with one to five substituents independently selected from the group consisting of phenyl, substituted phenyl, (C₁-C₆) alkyl and A, with the proviso that one and only one of said substituents is A, and one and only one of said substituents is phenyl or substituted phenyl, said substituted phenyl being substituted with one or more substituents independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) alkylthio, halogen and trifluoromethyl; and R⁴ is hydrogen, XR¹⁰ or A; with the proviso that when R¹ is a group of the formula XXVII or a group of the formula XXVI wherein G is nitrogen and wherein neither R⁵, R⁶ nor R¹⁵ is NR¹⁹R²⁰, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄) alkylthio, phenylthio or heteroalkylthio, R² and R³, together with the carbon to which they are attached, do not form a (C₃-C₇) cycloalkyl ring containing only carbon atoms:

A is a hydrocarbon containing 4 to 16 carbons and 0, 1 or 2 double bonds:

X is O, S, SO, SO₂, NH, NR²³CO or NSO₂R²⁴, wherein R²³ is hydrogen or (C₁-C₆) alkyl and R²⁴ is (C₁-C₆) alkyl, phenyl or (C₁-C₃) alkyl-phenyl:


R⁵, R⁶, R¹⁵ and R¹⁶ are each independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, iodo, (C₁-C₄)alkyl, (C₁-C₄) haloalkyl, (C₁-C₄) alkoxy, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄)alkylthio, substituted phenylthio, heteroarylthio, heteroaryloxy, and NR¹⁹R²⁰, wherein R¹⁹ and R²⁰ are the same or different and are selected from the group consisting of hydrogen, (C₁-C₄) alkyl, phenyl, substituted phenyl, (C₁-C₄) acyl, aroyl, and substituted aroyl, wherein said substituted phenyl and substituted aroyl groups are substituted with one or more substituents independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) alkylthio, halogen and trifluoromethyl or R¹⁹ and R²⁰, together with the nitrogen to which they are attached, form a piperidine or morpholine ring; and wherein R⁵, R⁶, R¹⁵ and R¹⁶, when attached to a bicyclic system, may be attached to either ring of such system, with the proviso that no more than 3 non-hydrogen substituents may be attached to any one ring of such system:

R⁷, R⁸ and R⁹ are the same or different; R⁷ is selected from the group consisting of hydrogen, (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, methyl and fluoro; and R⁸ and R⁹ are each independently selected from the group consisting of (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, methyl, and fluoro:

R¹⁰ is selected from the group consisting of (C₄-C₁₂) cycloalkyl, (C₄-C₁₂) straight or branched alkyl, (C₄-C₁₂) cycloalkyl-(C₁-C₆) alkyl, phenyl-(C₁-C₆) alkyl, (substituted phenyl)-(C₁-C₆) alkyl, (C₁-C₆) alkyl-phenyl, (C₁-C₆) alkyl-(substituted phenyl), substituted thiazoles, substituted benzothiazoles, and substituted pyridines; wherein the substituents on the substituted phenyl, substituted thiazoles, substituted benzothiazoles and substituted pyridines are selected from the group consisting of (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, (C₁-C₆) alkyl, halo and trifluoromethyl:


B, D, E and G are selected from the group consisting of nitrogen and carbon, with the proviso that one or more of B, D and E is nitrogen, and with the proviso that when G is nitrogen, the group XXVI is attached to the nitrogen of formula I at position 4 or 5 of the pyrimidine ring (designated by a and b):

or a pharmaceutically acceptable salt thereof, comprising:
reacting a compound of the formula




wherein W is chloro or bromo and R², R³ and R⁴ are defined as above, with an amine of the formula R¹NH₂, wherein R¹ is defined as above, and an acid scavenger.

2. A process according to claim 1, wherein said compound of the formula IV is obtained by reacting a compound of the formula




wherein R², R³ and R⁴ are defined as in claim 1, with a chlorinating or brominating agent.

3. A process for making a compound of the formula




IE

wherein


R¹ is

XXVI



XXIV

or



XXVII

R⁵, R⁶, R¹⁵ and R¹⁶ are each independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, iodo, (C₁-C₄)alkyl, (C₁-C₄) haloalkyl, (C₁-C₄) alkoxy, (C₁-C₆) alkylthio, (C₅-C₇) cycloalkylthio, phenyl (C₁-C₄)alkylthio, substituted phenylthio, heteroarylthio, heteroaryloxy, and NR¹⁹R²⁰, wherein R¹⁹ and R²⁰ are the same or different and are selected from the group consisting of hydrogen, (C₁-C₄) alkyl, phenyl, substituted phenyl, (C₁-C₄) acyl, aroyl, and substituted aroyl, wherein said substituted phenyl and substituted aroyl groups are substituted with one or more substituents independently selected from the group consisting of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) alkylthio, halogen and trifluoromethyl, or R¹⁹ and R²⁰, together with the nitrogen to which they are attached, form a piperidine or morpholine ring; and wherein R⁵, R⁶, R¹⁵ and R¹⁶, when attached to a bicyclic system, may be attached to either ring of such system, with the proviso that no more than 3 non-hydrogen substituents may be attached to any one ring of such system:

R⁷, R⁸ and R⁹ are the same or different; R⁷ is selected from the group consisting of hydrogen, (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, methyl and fluoro; and R⁸ and R⁹ are each independently selected from the group consisting of (C₁-C₄) alkoxy, (C₁-C₄) alkylthio, methyl, and fluoro:


B, D, E and G are selected from the group consisting of nitrogen and carbon, with the proviso that one or more of B, D and E is nitrogen, and with the proviso that when G is nitrogen, the group XXVI is attached to the nitrogen of formula IE at the 4 or 5 position of the pyrimidine ring (designated by a and b):

and R¹⁷ and R¹⁸ are each independently selected from the group consisting of (C₄-C₁₂) straight or branched alkyl, phenyl-(C₁-C₄) alkyl, and (C₁-C₆) alkylphenyl-(C₁-C₆) alkyl:

with the proviso that when R¹ is a group of the formula XXVII, R⁷, R⁸ and R⁹ are each methoxy: or pharmaceutically acceptable salt thereof,


comprising reacting a secondary amine of the formula NHR¹⁷R¹⁸, wherein R¹⁷ and R¹⁸ are as defined above, with a compound of the formula R¹NCO, wherein R¹ is as defined above.

4. A process for making a compound of the formula




10 wherein X and R¹⁰ are defined as in claim 1, and one of R³ and R⁴ is hydrogen and the other is selected from hydrogen, (C₁-C₄)alkyl or A, wherein A is a hydrocarbon containing 4 to 16 carbons and zero, one or two double bonds, comprising:

(a) reacting a compound of the formula



20

wherein R³ and R⁴ are defined as above, with a compound of the formula HXR¹⁰, wherein X and R¹⁰ are defined as above, and a base to produce a compound of the formula



30

and

(b) reacting said compound of formula VI' with a hydroxide compound.

5. A process according to claim 4, wherein said compound of the formula VI is obtained by a method comprising:

35 (a) reacting a compound of the formula




wherein R³ and R⁴ are defined as in claim 4, with thionyl chloride;

45 (b) reacting the product resulting from the reaction of step (a) with bromine in the presence of a catalytic amount of iodine; and

(c) reacting the product resulting from the reaction of step (b) with ethanol.

6. A process for making a compound of the formula




55


wherein R¹, X, R¹⁰, R³ and R⁴ are defined as in claim 4, comprising reacting a compound of the formula R¹NHCOCBrR³R⁴, wherein R¹, R³ and R⁴ are defined as above, with a compound of the formula HXR¹⁰,

wherein X and R¹⁰ are defined as above.


7. A process for making a compound of the formula



15 wherein R⁴ is hydrogen or A, R¹¹ is A, hydrogen, phenyl or substituted phenyl, A is a hydrocarbon containing 4 to 16 carbons and zero, one or two double bonds, and each of R¹² and R¹³ is independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl, or R¹² and R¹³, together with the carbons to which they are attached, form a benzene ring;
comprising conducting a Diels-Alder reaction between a compound of the formula




wherein R⁴ and R¹¹ are defined as above, and R¹¹ and the carboxyl group are trans to each other, and a diene of the formula



wherein R¹² and R¹³ are defined as above.

8. A process for making a compound of the formula




wherein


50 (a) R⁴ is (C₁-C₄) alkyl or A, and R² and R³ are each independently selected from the group consisting of hydrogen, (C₁-C₄) alkyl, A or XR¹⁰, wherein A, X and R¹⁰ are defined as in claim 1; or R² and R³, together with the carbon to which they are attached, form a cyclic or bicyclic system selected from the group consisting of (C₃-C₇) cycloalkyl, (C₆-C₁₄) bicycloalkyl, one or two carbons of said cycloalkyl and bicycloalkyl groups being optionally replaced with oxygen or sulfur; and aryl-fused and heteroaryl-fused systems containing 8 to 15 carbons atoms, one ring of said aryl-fused and heteroaryl-fused systems being aromatic and the ring containing the carbon to which R² and R³ are attached being non-aromatic, one of the carbons of said aromatic ring being optionally replaced by sulfur or oxygen, one or more carbons of said non-aromatic ring being optionally replaced by sulfur or oxygen, and one or more carbons of said aromatic ring being optionally replaced by nitrogen;

55


comprising reacting a compound of the formula



10 wherein R² and R³ are defined as above, with a base and then with a compound of the formula R⁴Hal, wherein Hal is halogen and R⁴ is (C₁-C₇)alkyl or A, wherein A is defined as above.
9. A process for preparing a compound of the formula



20 wherein R⁴ is XR¹⁰, wherein X and R¹⁰ are defined as in claim 1, and R² and R³, together with the carbons to which they are attached, form a cyclic or bicyclic system, wherein said cyclic or bicyclic system is defined as in claim 8;
comprising reacting a compound of the formula



wherein R² and R³ are defined as above, with a compound of the formula R¹⁰SSR¹⁰, wherein R¹⁰ is defined as above.

35 10. A process according to claim 2, wherein said acid scavenger is selected from the group consisting of dimethylaminopyridine, pyridine and triethylamine, and said chlorinating or brominating agent is selected from the group consisting of oxalyl chloride, oxalyl bromide, thionyl chloride, thionyl bromide, phosphorus trichloride, phosphorous tribromide, phosphorous pentachloride, phosphorous pentabromide, phosphorous oxychloride and phosphorous oxybromide.

40

45


50

55

10/616,579

ACCESSION NUMBER: 115:71632 CA
 TITLE: Preparation of new N-aryl and N-heteroaryl amide and
 -urea derivatives as inhibitors of acyl coenzyme
 A:cholesterol acyltransferase
 INVENTOR(S): McCarthy, Peter A.; Walker, Frederick J.; Truong,
 Thien; Hamanaka, Ernest S.; Chang, George
 PATENT ASSIGNEE(S): Pfizer Inc., USA
 SOURCE: Eur. Pat. Appl., 85 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 418071 ✓	A2	19910320	EP 1990-310009	19900913
EP 418071 ✓	A3	19920325		
EP 418071 ✓	B1	19950426		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
WO 9104027	A1	19910404	WO 1989-US4033	19890915
W: FI, HU, NO, RO, SU, US				
IL 95610	A1	19941229	IL 1990-95610	19900907
DD 298092	A5	19920206	DD 1990-343971	19900912
CA 2025301	AA	19910316	CA 1990-2025301	19900913
CA 2025301	C	20011016		
EP 609960	A1	19940810	EP 1994-200437	19900913
EP 609960	B1	19990303		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
AT 121730	E	19950515	AT 1990-310009	19900913
ES 2071033	T3	19950616	ES 1990-310009	19900913
AT 177082	E	19990315	AT 1994-200437	19900913
ES 2127878	T3	19990501	ES 1994-200437	19900913
NO 9004022	A	19910318	NO 1990-4022	19900914
CN 1050183	A	19910327	CN 1990-108294	19900914
HU 54625	A2	19910328	HU 1990-5991	19900914
AU 9062553	A1	19910418	AU 1990-62553	19900914
AU 652345	B2	19940825		
JP 03120243	A2	19910522	JP 1990-245969	19900914
JP 08025974	B4	19960313		
ZA 9007346	A	19920527	ZA 1990-7346	19900914
PL 165370	B1	19941230	PL 1990-286899	19900914
PL 165357	B1	19941230	PL 1990-291470	19900914
HU 70027	A2	19950928	HU 1993-2945	19900914
FI 111362	B1	20030715	FI 1990-4537	19900914
PRIORITY APPLN. INFO.:			WO 1989-US4033	A 19890915
			EP 1990-310009	A3 19900913
OTHER SOURCE(S):	MARPAT 115:71632			
GI				



I

10/616,579

AB Approx. 250 title amides and ureas R_1NHCOQ [$Q = CR_2R_3R_4$, NR_5R_6 ; $R_1 =$ (substituted) pyridyl, pyrimidinyl, quinolinyl, pyridoimidazolyl, etc., substituted Ph; $R_2-R_4 = H$, alkyl, hydrocarbyl, XR_7 , phenylalkyl, cycloalkyl; or R_3R_4 forms cycloalkyl, cycloalkenyl, bicycloalkyl, etc.; $R_5, R_6 =$ alkyl, phenylalkyl, alkylphenylalkyl; $R_7 =$ alkyl, cycloalkyl, phenylalkyl, thiazolyl, pyridyl, etc.; $X = O, S, SO, SO_2, NH$, etc.; numerous provisos] were prepared as hypolipidemics and antiatherosclerotics (no data). For example, 2-(hexylthio)decanoic acid was refluxed with $SOCl_2$ in C_6H_6 to give the acid chloride, which was added to 5-amino-4,6-bis(methylthio)-2-methylpyrimidine in CH_2Cl_2 followed by refluxing and purification to give 72.4% title amide I.

IT 134989-87-0P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of, as hypolipidemic)

RN 134989-87-0 CA

CN Urea, N-heptyl-N'-5-isoquinolinyl-N-[[4-(3-methylbutyl)phenyl]methyl]-
(9CI) (CA INDEX NAME)

